

**INVITRO AND INVIVO EVALUATION OF ANTICANCER
ACTIVITY OF DIFFERENT EXTRACTS OF *IPOMOEA*
BATAUS STEM**

A Dissertation submitted to
THE TAMIL NADU Dr. M.G.R. MEDICAL UNIVERSITY,
CHENNAI- 600 032

In partial fulfillment of the requirements for the award of the Degree of
MASTER OF PHARMACY
IN
BRANCH -VI - PHARMACOLOGY

Submitted by
Ms. ANOOPA WILSON
REGISTRATION No.261526151

Under the guidance of
Prof. G.NAGARAJA PERUMAL, M.Pharm., (Ph.D).,
Professor & Head
Department of Pharmacology



DEPARTMENT OF PHARMACOLOGY
KARPAGAM COLLEGE OF PHARMACY
COIMBATORE-641 032

MAY - 2017

CERTIFICATE

This is to certify that the M.Pharm Dissertation entitled **“Invitro And *Invivo* Evaluation Of Different Extract Of *ipomoea batatus* stem”** being submitted to The TamilNadu Dr. M.G.R Medical University, Chennai was carried out by **Ms. ANOOPA WILSON** to The Tamil Nadu Dr. M.G.R Medical University, Chennai in partial fulfillment for the degree of **MASTER OF PHARMACY IN PHARMACOLOGY** is a bonafied work carried out by candidate under my guidance and supervision in the Department of Pharmacology , Karpagam college of Pharmacy Coimbatore – 32.

I have fully satisfied with her performance and work. I have forwarded this dissertation work for evaluation.

Station :

G.NAGARAJA PERUMAL M Pharm.,(Ph.D).,

Date :

Professor & Head

Department of Pharmacology

CERTIFICATE

This is to certify that the M.Pharm Dissertation entitled **“*Invitro* And *Invivo* Evaluation Of Different Extracts Of *Ipomoea batatus* stem”** being submitted to The Tamil Nadu Dr. M.G.R Medical University, Chennai was carried out by **Ms. ANOOPA WILSON** to The Tamil Nadu Dr.M.G.R Medical University , Chennai in partial fulfillment for the degree of **MASTER OF PHARMACY IN PHARMACOLOGY** is a bonafied work carried out by candidate under the guidance of **Prof. G. Nagaraja Perumal, M.Pharm., (Ph.D).**, in the Department of Pharmacology, Karpagam college of Pharmacy Coimbatore – 32.

I have fully satisfied with her performance and work. I have forwarded this dissertation work for evaluation.

Station :

Dr .S.MOHAN M.Pharm.,Ph.D.

Date: :

Principal

DECLARATION

I hereby declare that this dissertation "***Invitro And Invivo*** Evaluation Of Different Extracts of ***Ipomoea Batatus stem***" submitted by the candidate, in partial fulfillment of requirements for the degree of **MASTER OF PHARMACY IN PHARMACOLOGY** to The Tamil Nadu Dr.M.G.R Medical University, Chennai is the result of my original and independent research work carried out under the guidance of **Prof .G.NAGARAJA PERUMAL., M.Pharm.,(Ph.D)** Professor & Head Department of Pharmacology ,Karpagam College of Pharmacy, Coimbatore -32,& Co Guide **Dr Hashim K.M.,UWIN LIFE SCIENCE**, during the academic year 2016-2017.

Station :

ANOOPA WILSON

Date :

Reg . No . 261526151

EVALUATION CERTIFICATE

This is to certify that disseration work entitled “ ***Invitro And Invivo*** **Evaluation Of Different Extracts Of *Ipomoea Batatus* Stem ”** submitted by **Ms. Anoop Wilson**, bearing **Reg. No : 261526151** to the The Tamil Nadu Dr.M.G.R Medical University, Chennai in the partial fulfillment for the degree of **MASTER OF PHARMACY IN PHARMACOLOGY** is a bonafied work carried out during the academic year 2016-2017 by the candidate at **Department of Pharmacology**, Karpagam College of Pharmacy, Coimbatore and evaluated by us.

Examination centre :

Date :

Internal Examiner

Convenor of Examination

External examiner

**DEDICATED TO MY BELOVED
PARENTS, TEACHERS AND
ALMIGHTY**

ACKNOWLEDGEMENT

ACKNOWLEDGEMENT

First of all I would like to thank god for his blessings to do this research work successfully. With immense pleasure and pride, I would like to take this opportunity in expressing my deep sense of gratitude to my beloved guide **Mr G.Nagaraja Perumal M.Pharm.**, Professor & Head, Department of Pharmacology, Karpagam College of Pharmacy under whose active guidance, innovative ideas, constant inspiration and encouragement of the work entitled “***Invitro And Invivo Evaluation Of Different Extracts Of Ipomoea Batatus Stem***” has been carried out.

I wish to express my deep sense of gratitude to Dr.R.Vasanthakumar Chairman of Karpagam Group of institutions for the facilities provided me in this institution.

My sincere thanks to our respected and beloved Principal Dr.S.Mohan, M Pharm ,Ph.D, Karpagam College of Pharmacy for his encouragement and also providing all facilities in this institution to the fullest possible extent enabling me to complete this work successfully.

I convey my gratitude to Mrs.V.Idachristi M. Pharm,Professor & Head ,Department of Pharmacognosy helped me to proceed useful ideas.

My whole hearted thanks to Mr.D. Ranjith kumar M Pharm,Asst. Professor,Department of Pharmaceutical Analysis for his kind advice.

I am also conveying my thanks to Mrs. M. Karpagavalli ,M. Pharm, Associate Professor, Department of Pharmaceutical chemistry, for encouragement and valuable suggestion during this work.

I express my sincere thanks to Mr. K. Nahas , Lab assistant , Department of Pharmaceutical chemistry for his kind support.

I convey my gratitude to Mr. S. Asker , Lab Assistant , Department of Pathology for his kind support.

I express my sincere thanks to Mrs.M. Sathybhama Lab assistant, Department of Pharmaceutical chemistry for her kind support.

I am duly bound to all my non teaching staffs of Karpagam college of Pharmacy for their valuable advices and co-operation. Above all , I am remain indebted to my seniors class mates (mohammed shanavas v.k Bhavan, Shanavas, Amritha, Habeeb, Sijad, Ubaid), to my beloved parents who inspired and guided me and also for being the backbone for all my successful endeavors in my life.

ANOOPA WILSON
(261526151)

CONTENTS

SL NO	CONTENT	PAGE NO
1	INTRODUCTION	1
2	REVIEW OF LITERATURE	24
3	AIM AND OBJECTIVE	35
4	PLAN OF WORK	36
5	PLANT PROFILE	37
6	MATERIALS AND METHODS	41
7	RESULTS AND DISCUSSION	49
8	SUMMARY AND CONCLUSION	61
9	BIBLIOGRAPHY	62

CHAPTER I

1.INTRODUCTION

1.1 CANCER

Cancer is a diseases involving abnormal cell growth with the potential to invade or spread to other parts of the body.^{[1]&[2]} Not all tumors are cancerous; benign tumors do not spread to other parts of the body.^[2] Possible signs and symptoms include a lump, abnormal bleeding, prolonged cough, unexplained weight loss and a change in bowel movements. While these symptoms may indicate cancer, they may have other causes.^[3] Over 100 cancers affect humans.^[2]

Many cancers can be prevented by not smoking, maintaining a healthy weight, not drinking too much alcohol, eating plenty of vegetables, fruits and whole grains, vaccination against certain infectious diseases, not eating too much processed and red meat, and avoiding too much sunlight exposure.^{[9]&[10]} Early detection through screening is useful for cervical and colorectal cancer.^[11] The benefits of screening in breast cancer are controversial.^{[11]&[12]} Cancer is often treated with some combination of radiation therapy, surgery, chemotherapy, and targeted therapy.^{[1]&[13]} Pain and symptom management are an important part of care. Palliative care is particularly important in people with advanced disease.^[1] The chance of survival depends on the type of cancer and extent of disease at the start of treatment.^[6] In children under 15 at diagnosis the five-year survival rate in the developed world is on average 80%.^[14] For cancer in the United States the average five-year survival rate is 66%.^[15]

Worldwide approximately 18% of cancer deaths are related to infectious diseases. This proportion ranges from a high of 25% in Africa to less than 10% in the developed world.^[5] Viruses are the

usual infectious agents that cause cancer but cancer bacteria and parasites may also play a role.

In 2012 about 14.1 million new cases of cancer occurred globally (not including skin cancer other than melanoma).^[6] It caused about 8.2 million deaths or 14.6% of human deaths.^{[6]&[16]} The most common types of cancer in males are lung cancer, prostate cancer, colorectal cancer and stomach cancer. In females, the most common types are breast cancer, colorectal cancer, lung cancer and cervical cancer.^[6] If skin cancer other than melanoma were included in total new cancers each year it would account for around 40% of cases.^{[17]&[18]} In children, acute lymphoblastic leukaemia and brain tumors are most common except in Africa where non-Hodgkin lymphoma occurs more often.^[14] In 2012, about 165,000 children under 15 years of age were diagnosed with cancer. The risk of cancer increases significantly with age and many cancers occur more commonly in developed countries.^[6] Rates are increasing as more people live to an old age and as lifestyle changes occur in the developing world.^[19] The financial costs of cancer were estimated at \$1.16 trillion US dollars per year as of 2010.^[20]

Cancers are a large family of diseases that involve abnormal cell growth with the potential to invade or spread to other parts of the body.^{[1]&[2]} They form a subset of neoplasms. A neoplasm or tumor is a group of cells that have undergone unregulated growth and will often form a mass or lump, but may be distributed diffusely.^{[21][22]}

All tumor cells show the six hallmarks of cancer. These characteristics are required to produce a malignant tumor. They include:^[23]

- Cell growth and division absent the proper signals
- Continuous growth and division even given contrary signals
- Avoidance of programmed cell death
- Limitless number of cell divisions

- Promoting blood vessel construction
- Invasion of tissue and formation of metastases[24]

The progression from normal cells to cells that can form a detectable mass to outright cancer involves multiple steps known as malignant progression.^{[24]&[25]}

When cancer begins, it produces no symptoms. Signs and symptoms appear as the mass grows or ulcerates. The findings that result depend on the cancer's type and location. Few symptoms are specific. Many frequently occur in individuals who have other conditions. Cancer is a "great imitator". Thus, it is common for people diagnosed with cancer to have been treated for other diseases, which were hypothesized to be causing their symptoms.^[26]

People may become anxious or depressed post-diagnosis. The risk of suicide in people with cancer is approximately double.^[27]

Local symptoms may occur due to the mass of the tumor or its ulceration. For example, mass effects from lung cancer can block the bronchus resulting in cough or pneumonia; esophageal cancer can cause narrowing of the esophagus, making it difficult or painful to swallow; and colorectal cancer may lead to narrowing or blockages in the bowel, affecting bowel habits. Masses in breasts or testicles may produce observable lumps. Ulceration can cause bleeding that, if it occurs in the lung, will lead to coughing up blood, in the bowels to anemia or rectal bleeding, in the bladder to blood in the urine and in the uterus to vaginal bleeding. Although localized pain may occur in advanced cancer, the initial swelling is usually painless. Some cancers can cause a buildup of fluid within the chest or abdomen.^[26]

1.1.2 Systemic symptoms

General symptoms occur due to effects that are not related to direct or metastatic spread. These may include: unintentional

weight loss, fever.^[28] Hodgkin disease, leukemia and cancers of the liver or kidney can cause a persistent fever.^[26]

Some cancers may cause specific groups of systemic symptoms, termed paraneoplastic phenomena.^[26]

1.1.2.3 Mechanism Of Cancer Cell Formation And Spreading

Cancer can spread from its original site by local spread, lymphatic spread to regional lymph nodes or by hematogenous spread via the blood to distant sites, known as metastasis. When cancer spreads by a hematogenous route, it usually spreads all over the body. However, cancer 'seeds' grow in certain selected site only ('soil') as hypothesized in the soil and seed hypothesis of cancer metastasis. The symptoms of metastatic cancers depend on the tumor location and can include enlarged lymph nodes (which can be felt or sometimes seen under the skin and are typically hard), enlarged liver or enlarged spleen, which can be felt in the abdomen, pain or fracture of affected bones and neurological symptoms.^[26]

1.2 Causes

The majority of cancers, some 90–95% of cases, are due to environmental factors. The remaining 5–10% are due to inherited genetics. Environmental, as used by cancer researchers, means any cause that is not inherited genetically, such as lifestyle, economic and behavioral factors and not merely pollution.^[29] Common environmental factors that contribute to cancer death include tobacco (25–30%), diet and obesity (30–35%), infections (15–20%), radiation, stress.


Tobacco use is the cause of about 22% of cancer deaths.^[1] Another 10% is due to obesity, poor diet, lack of physical activity and drinking alcohol.^{[1]&[4]} Other factors include certain infections,

exposure to ionizing radiation and environmental pollutants.^[5] It is due to infections such as hepatitis B, hepatitis C and human papillomavirus (HPV).^[1] These factors act, at least partly, by changing the genes of a cell. Typically many genetic changes are required before cancer develops.^[6] Approximately 5–10% of cancers are due to inherited genetic defects from a person's parents.^[7] Cancer can be detected by certain signs and symptoms or screening tests.^[1] It is then typically further investigated by medical imaging and confirmed by biopsy.^[8]

It is not generally possible to prove what caused a particular cancer because the various causes do not have specific fingerprints. For example, if a person who uses tobacco heavily develops lung cancer, then it was probably caused by the tobacco use, but since everyone has a small chance of developing lung cancer as a result of air pollution or radiation, the cancer may have developed for one of those reasons. Excepting the rare transmissions that occur with pregnancies and occasional organ donors, cancer is generally not a transmissible disease.^[30]

1.21 Chemicals

Figure No.1: Lag time between Smoking and Lung Cancer



The incidence of lung cancer is highly correlated with smoking.

Exposure to particular substances have been linked to specific types of cancer. These substances are called carcinogens.

Tobacco smoke, for example, causes 90% of lung cancer.^[31] It also causes cancer in the larynx, head, neck, stomach, bladder, kidney, esophagus and pancreas.^[32] Tobacco smoke contains over fifty known carcinogens, including nitrosamines and polycyclic aromatic hydrocarbons.^[33]

Tobacco is responsible about one in five cancer deaths worldwide^[33] and about one in three in the developed world.^[34] Lung cancer death rates in US have mirrored smoking patterns, with increases in smoking followed by dramatic increases in lung cancer death rates and, more recently, decreases in smoking rates since the 1950s followed by decreases in lung cancer death rates in men since 1990.^{[35]&[36]}

In Western Europe, 10% of cancers in males and 3% of cancers in females are attributed to alcohol exposure, especially liver and digestive tract cancers.^[37] Cancer from work-related substance exposures may cause between 2 and 20% of cases,^[38] causing at least 200,000 deaths. Cancers such as lung cancer and mesothelioma can come from inhaling tobacco smoke or asbestos fibers, or leukemia from exposure to benzene^[39]

1.2.2 Life style Changes

Diet, physical inactivity and obesity are related to up to 30–35% of cancer deaths.^{[5][40]} In the United States excess body weight is associated with the development of many types of cancer and is a factor in 14–20% of cancer deaths.^[40] A UK study including data on over 5 million people showed higher body mass index to be related to at least 10 types of cancer and responsible for around 12,000

cases each year in that country.^[41] Physical inactivity is believed to contribute to cancer risk, not only through its effect on body weight and also through negative effects on the immune system and endocrine system. More than half of the effect from diet is due to overnutrition (eating too much), rather than from eating too few vegetables or other healthful foods.

Some specific foods are linked to specific cancers. A high-salt diet is linked to gastric cancer.^[42] Aflatoxin B₁, a frequent food contaminant, causes liver cancer. Betel nut chewing can cause oral cancer.^[42] National differences in dietary practices may partly explain differences in cancer incidence. For example, gastric cancer is more common in Japan due to its high-salt diet while colon cancer is more common in the United States. Immigrant cancer profiles develop mirror that of their new country, often within one generation.^[44]

1.2.3. Infection

Oncoviruses (viruses that can cause cancer) include human papillomavirus (cervical cancer), Epstein–Barr virus (B-cell lymphoproliferative disease and nasopharyngeal carcinoma), Kaposi's sarcoma herpesvirus (Kaposi's sarcoma and primary effusion lymphomas), hepatitis B and hepatitis C viruses (hepatocellular carcinoma) and human T-cell leukemia virus-1 (T-cell leukemias).

1.2.4. Radiation

Up to 10% of invasive cancers are related to radiation exposure, including both ionizing radiation and non-ionizing ultraviolet radiation.^[5] Additionally, the majority of non-invasive cancers are non-melanoma skin cancers caused by non-ionizing ultraviolet radiation, mostly from sunlight. Sources of ionizing radiation include medical imaging and radon gas.

Ionizing radiation is not a particularly strong mutagen.[48] Residential exposure to radon gas, for example, has similar cancer risks as passive smoking. Radiation is a more potent source of cancer when combined with other cancer-causing agents, such as radon plus tobacco smoke. Radiation can cause cancer in most parts of the body, in all animals and at any age..

Medical use of ionizing radiation have small growing source of radiation-induced cancers. Ionizing radiation may be used to treat other cancers, but this may, in some cases, induce a second form of cancer.^[48] It is also used in some kinds of medical imaging.^[49]

Prolonged exposure to ultraviolet radiation from the sun can lead to melanoma and other skin malignancies. Clear evidence establishes ultraviolet radiation, especially the non-ionizing medium wave UVB, as the cause of most non-melanoma skin cancers, which are the most common forms of cancer in the world.^[50]

Non-ionizing radio frequency radiation from mobile phones, electric power transmission and other similar sources have been described as a possible carcinogen by the World Health Organization's International Agency for Research on Cancer.^[51] However, studies have not found a consistent link between mobile phone radiation and cancer risk.^[52]

Some substances cause cancer primarily through their chemical rather than physical, effects. A prominent example of this is prolonged exposure to asbestos, naturally occurring mineral fibers that are a major cause of mesothelioma (cancer of the serous membrane) usually the serous membrane surrounding the lungs. Other substances in this category, including both naturally occurring and synthetic asbestos-like fibers, such as wollastonite, attapulgite, glass wool and rock wool, are believed metallic cobalt and nickel and crystalline silica (quartz, cristobalite

and tridymite). Usually, physical carcinogens must get inside the body (such as through inhalation) and require years of exposure to produce cancer.[55] Children and adolescents are twice as likely to develop radiation-induced leukemia as adults; radiation exposure before birth has ten times the effect

Physical trauma resulting in cancer is relatively rare. Claims that breaking bones resulted in bone cancer, for example, have not been proven. Similarly, physical trauma is not accepted as a cause for cervical cancer, breast cancer or brain cancer. One accepted source is frequent, long-term application of hot objects to the body. It is possible that repeated burns on the same part of the body, such as those produced by kanger and kairo heaters (charcoal hand warmers), may produce skin cancer, especially if carcinogenic chemicals are also present. Frequent consumption of scalding hot tea may produce esophageal cancer.^[56] Generally, it is believed that cancer arises, or a pre-existing cancer is encouraged, during the process of healing, rather than directly by the trauma. However, repeated injuries to the same tissues might promote excessive cell proliferation, which could then increase the odds of a cancerous mutation.^[56]

Chronic inflammation has been hypothesized to directly cause mutation.^{[56][57]} Inflammation can contribute to proliferation, survival, angiogenesis and migration of cancer cells by influencing the tumor microenvironment.^{[58]&[59]} Oncogenes build up an inflammatory pro-tumorigenic microenvironment.^[60]

1.2.5 Hormones

Some hormones play a role in the development of cancer by promoting cell proliferation.^[61] Insulin-like growth factors and their binding proteins play a key role in cancer cell proliferation, differentiation, apoptosis, suggesting possible involvement in carcinogenesis.^[62]

Hormones are important agents in sex-related cancers, such as cancer of the breast, endometrium, prostate, ovary, testis and also of thyroid cancer and bone cancer. For example, the daughters of women who have breast cancer have significantly higher levels of estrogen and progesterone than the daughters of women without breast cancer. These higher hormone levels may explain their higher risk of breast cancer, even in the absence of a breast-cancer gene. Similarly, men of African ancestry have significantly higher levels of testosterone than men of European ancestry and have a correspondingly higher level of prostate cancer. Men of Asian ancestry, with the lowest levels of testosterone-activating androstenediolglucuronide, have the lowest levels of prostate cancer.^[61]

Other factors are relevant: obese people have higher levels of some hormones associated with cancer and a higher rate of those cancers. Women who take hormone replacement therapy have higher risk of developing cancers associated with those hormones. On the other hand, people who exercise far more than average have lower levels of these hormones and lower risk of cancer. Osteosarcoma may be promoted by growth hormones. Some treatments and prevention approaches leverage this cause by artificially reducing hormone levels and thus discouraging hormone-sensitive cancers.^[61] Cancer is fundamentally disease of tissue growth regulation. In order for a normal cell to transform into a cancer cell, the genes that regulate cell growth and differentiation must be altered.^[65]

The affected genes are divided into two broad categories. Oncogenes are genes that promote cell growth and reproduction. Tumor suppressor genes are genes that inhibit cell division and survival. Malignant transformation can occur through the formation of novel oncogenes, the inappropriate over-expression of normal oncogenes, or by the under-expression or disabling of tumor suppressor genes. Typically, changes in multiple

genes are required to transform a normal cell into a cancer cell.[66]

Genetic changes can occur by different mechanisms. The gain or loss of an entire chromosome can occur through errors in mitosis. More common are mutations, which are changes in the nucleotide sequence of genomic DNA.

Large-scale mutations involve the deletion or gain of a portion of a chromosome. Genomic amplification occurs when a cell gains copies (often 20 or more) of a small chromosomal locus, usually containing one or more oncogenes and adjacent genetic material. Translocation occurs when two separate chromosomal regions become abnormally fused, often at a characteristic location. A well-known example of this is the Philadelphia chromosome, or translocation of chromosomes 9 and 22, which occurs in chronic myelogenous leukemia and results in production of the BCR-abl fusion protein, an oncogenic tyrosine kinase.

Small-scale mutations include point mutations, deletions, and insertions, which are in the promoter region of a gene and affect its expression, or may occur in the gene's coding sequence and alter the function or stability of its protein product. Disruption of a single gene may also result from integration of genomic material from a DNA virus or retrovirus, leading to the expression of viral oncogenes in the affected cell and its descendants.

Replication of the data contained within the DNA of living cells will probabilistically result in some errors (mutations). Complex error correction and prevention is built into the process and safeguards the cell against cancer. If a significant error occurs, the damaged cell can self-destruct through programmed cell death, termed apoptosis. If the error control processes fail, then the mutations will survive and be passed along to daughter cells.

Some environments make errors more likely to arise and propagate. Such environments can include the presence of


disruptive substances called carcinogens, repeated physical injury, heat, ionising radiation or hypoxia.^[67]

The errors that cause cancer are self-amplifying and compounding, for example:

- A mutation in the error-correcting machinery of a cell might cause that cell and its children to accumulate errors more rapidly.
- A further mutation in an oncogene might cause the cell to reproduce more rapidly and more frequently than its normal counterparts.
- A further mutation may cause loss of a tumor suppressor gene, disrupting the apoptosis signaling pathway and immortalizing the cell.
- A further mutation in the signaling machinery of the cell might send error-causing signals to nearby cells.

The transformation of a normal cell into cancer is akin to a chain reaction caused by initial errors, which compound into more severe errors, each progressively allowing the cell to escape more controls that limit normal tissue growth. This rebellion-like scenario is an undesirable survival of the fittest, where the driving forces of evolution work against the body's design and enforcement of order. Once cancer has begun to develop, this ongoing process, termed clonal evolution, drives progression towards more invasive stages.^[68] Clonal evolution leads to intra-tumour heterogeneity (cancer cells with heterogeneous mutations) that complicates designing effective treatment strategies

Figure No:2 Cell Proliferation



Characteristic abilities developed by cancers are divided into categories, specifically evasion of apoptosis, self-sufficiency in growth signals, insensitivity to anti-growth signals, sustained angiogenesis, limitless replicative potential, metastasis, reprogramming of energy metabolism and evasion of immune destruction.^{[24]&[25]}


The classical view of cancer is a set of diseases that are driven by progressive genetic abnormalities that include mutations in tumor-suppressor genes and oncogenes and chromosomal abnormalities. Later epigenetic alterations' role was identified.^[69]

Epigenetic alterations refer to functionally relevant modifications to the genome that do not change the nucleotide sequence. Examples of such modifications are changes in DNA methylation (hypermethylation and hypomethylation), histone

modification[70] and changes in chromosomal architecture (caused by inappropriate expression of proteins such as HMGA2 or HMGA1).^[71] Each of these alterations regulates gene expression without altering the underlying DNA sequence. These changes may remain through cell divisions, last for multiple generations and can be considered to be epimutations (equivalent to mutations).

Epigenetic alterations occur frequently in cancers. As an example, one study listed protein coding genes that were frequently altered in their methylation in association with colon cancer. These included 147 hypermethylated and 27 hypomethylated genes. Of the hypermethylated genes, 10 were hypermethylated in 100% of colon cancers and many others were hypermethylated in more than 50% of colon cancers.^[72]

Figure No:3



While epigenetic alterations are found in cancers, the epigenetic alterations in DNA repair genes, causing reduced expression of DNA repair proteins, may be of particular importance. Such alterations are thought to occur early in progression to cancer and to be a likely cause of the genetic instability characteristic of cancers.^{[73],[74],[75]&[76]}

Reduced expression of DNA repair genes disrupts DNA repair. This is shown in the figure at the 4th level from the top. (In the figure, red wording indicates the central role of DNA damage and defects in DNA repair in progression to cancer.) When DNA repair is deficient DNA damage remains in cells at a higher than usual level (5th level) and cause increased frequencies of mutation and/or epimutation (6th level). Mutation rates increase substantially in cells defective in DNA mismatch repair^{[77]&[78]} or in homologous recombinational repair (HRR).^[79] Chromosomal rearrangements and aneuploidy also increase in HRR defective cells.^[80]

Higher levels of DNA damage cause increased mutation (right side of figure) and increased epimutation. During repair of DNA double strand breaks, or repair of other DNA damage, incompletely cleared repair sites can cause epigenetic gene silencing.^{[81][82]}

Deficient expression of DNA repair proteins due to an inherited mutation can increase cancer risks. Individuals with an inherited impairment in any of 34 DNA repair genes (see article DNA repair-deficiency disorder) have increased cancer risk, with some defects ensuring a 100% lifetime chance of cancer (e.g. p53 mutations).^[83] Germ line DNA repair mutations are noted on the figure's left side. However, such germline mutations (which cause highly penetrant cancer syndromes) are the cause of only about 1 percent of cancers.^[84]

In sporadic cancers, deficiencies in DNA repair are occasionally caused by a mutation in a DNA repair gene but are much more frequently caused by epigenetic alterations that reduce or silence

expression of DNA repair genes. This is indicated in the figure at the 3rd level. Many studies of heavy metal-induced carcinogenesis show that such heavy metals cause a reduction in expression of DNA repair enzymes, some through epigenetic mechanisms. DNA repair inhibition is proposed to be a predominant mechanism in heavy metal-induced carcinogenicity. In addition, frequent epigenetic alterations of the DNA sequences code for small RNAs called microRNAs (or miRNAs). miRNAs do not code for proteins, but can "target" protein-coding genes and reduce their expression.

Cancers usually arise from an assemblage of mutations and epimutations that confer a selective advantage leading to clonal expansion.^[85]

Cancer is becoming a high profile disease in developed and developing worlds. In 2007 the WHO published that in 2005, 7.6 million people died from cancer related diseases with the majority of these people living in low-income countries . In the United States cancer is the cause of 1 in 4 deaths and in 2010 it was estimated there were over 1.5 million new cases of cancer . Cancer Research UK said in 2012 14.1 million adults were diagnosed with cancer and 8.2 million people were killed by cancer globally . Therefore, the demand for a cure and the prevention of cancer is extremely high.

1.2.6. Dietary

While many dietary recommendations have been proposed to reduce cancer risks, the evidence to support them is not definitive.^{[9]&[93]} The primary dietary factors that increase risk are obesity and alcohol consumption. Diets low in fruits and vegetables and high in red meat have been implicated but reviews and meta-analyses do not come to a consistent conclusion.^{[94]&[95]}

A 2014 meta-analysis find no relationship between fruits and vegetables and cancer.^[96] Coffee is associated with a reduced risk

of liver cancer.^[97] Studies have linked excess consumption of red or processed meat to an increased risk of breast cancer, colon cancer and pancreatic cancer, a phenomenon that could be due to the presence of carcinogens in meats cooked at high temperatures.^{[98]&[99]} In 2015 the IARC reported that eating processed meat (e.g., bacon, ham, hot dogs, sausages) and, to a lesser degree, red meat was linked to some cancers.^{[100]&[101]} Dietary recommendations for cancer prevention typically include an emphasis on vegetables, fruit, whole grains and fish and an avoidance of processed and red meat (beef, pork, lamb), animal fats and refined carbohydrates.^{[9]&[93]}

1.2.7. Medication

Medications can be used to prevent cancer in a few circumstances.^[102] In the general population, NSAIDs reduce the risk of colorectal cancer; however, due to cardiovascular and gastrointestinal side effects, they cause overall harm when used for prevention.^[103] Aspirin has been found to reduce the risk of death from cancer by about 7%.^[104] COX-2 inhibitors may decrease the rate of polyp formation in people with familial adenomatous polyposis; however, it is associated with the same adverse effects as NSAIDs.^[105] Daily use of tamoxifen or raloxifene reduce the risk of breast cancer in high-risk women.^[106] The benefit versus harm for 5-alpha-reductase inhibitor such as finasteride is not clear.^[107]

Vitamins are not effective at preventing cancer,^[108] although low blood levels of vitamin D are correlated with increased cancer risk.^{[109]&[110]} People who have cancer are also at a high risk of developing vitamin D deficiency.^[111] Whether this relationship is causal and vitamin D supplementation is protective is not determined.^[112] Beta-carotene supplementation increases lung cancer rates in those who are high risk.^[113] Folic acid supplementation is not effective in preventing colon cancer

and may increase colon polyps.^[114] It is unclear if selenium supplementation has an effect.^[115]

1.2.8. Vaccination

Vaccines have been developed that prevent infection by some carcinogenic viruses. Human papillomavirus vaccine (Gardasil and Cervarix) decrease the risk of developing cervical cancer. The hepatitis B vaccine prevents infection with hepatitis B virus and thus decreases the risk of liver cancer.^[116] The administration of human papillomavirus and hepatitis B vaccinations is recommended when resources allow.^[117]

1.2.9. Screening

Unlike diagnostic efforts prompted by symptoms and medical signs, cancer screening involves efforts to detect cancer after it has formed, but before any noticeable symptoms appear. This may involve physical examination, blood or urine tests or medical imaging.^[118]

Cancer screening is not available for many types of cancers. Even when tests are available, they may not be recommended for everyone. Universal screening or mass screening involves screening everyone.^[119] Selective screening identifies people who are at higher risk, such as people with a family history. Several factors are considered to determine whether the benefits of screening outweigh the risks and the costs of screening.^[118] These factors include:

- Possible harms from the screening test: for example, X-ray images involve exposure to potentially harmful ionizing radiation
- The likelihood of the test correctly identifying cancer
- The likelihood that cancer is present: Screening is not normally useful for rare cancers.

- Possible harms from follow-up procedures
- Whether suitable treatment is available
- Whether early detection improves treatment outcomes
- Whether the cancer will ever need treatment
- Whether the test is acceptable to the people: If a screening test is too burdensome (for example, extremely painful), then people will refuse to participate.^[119]
- Cost

1.3 Diagnosis

Unlike diagnostic efforts prompted by symptoms and medical signs, cancer screening involves efforts to detect cancer after it has formed, but before any noticeable symptoms appear. This may involve physical examination, blood or urine tests or medical imaging.

Cancer screening is not available for many types of cancers. Even when tests are available, they may not be recommended for everyone. Universal screening or mass screening involves screening everyone. Selective screening identifies people who are at higher risk, such as people with a family history. Several factors are considered to determine whether the benefits of screening outweigh the risks and the costs of screening. These factors include:

- Possible harms from the screening test: for example, X-ray images involve exposure to potentially harmful ionizing radiation
- The likelihood of the test correctly identifying cancer
- The likelihood that cancer is present: Screening is not normally useful for rare cancers.
- Possible harms from follow-up procedures
- Whether suitable treatment is available
- Whether early detection improves treatment outcomes

- Whether the cancer will ever need treatment
- Whether the test is acceptable to the people: If a screening test is too burdensome (for example, extremely painful), then people will refuse to participate

Chemically-derived drugs have been developed and other cancer treatments pre-exist. However, current methods such as chemotherapy have their limitations due to their toxic effects on non-targeted tissues furthering human health problems. Therefore, there is a demand for alternative treatments with naturally-derived anticancer agents with plants being the desired source.

1.4 Prevention

Cancer prevention is defined as active measures to decrease cancer risk. The vast majority of cancer cases are due to environmental risk factors. Many of these environmental factors are controllable lifestyle choices. Thus, cancer is generally preventable. Between 70% and 90% of common cancers are due to environmental factors and therefore potentially preventable.^[91]

Greater than 30% of cancer deaths could be prevented by avoiding risk factors including: tobacco, excess weight/obesity, insufficient diet, physical inactivity, alcohol, sexually transmitted infections and air pollution. Not all environmental causes are controllable, such as naturally occurring background radiation and cancers caused through hereditary genetic disorders and thus are not preventable via personal behavior.

1.5. Classification

Cancers are classified by the type of cell that the tumor cells resemble and is therefore presumed to be the origin of the tumor. These types include:

- **Carcinoma:** Cancers derived from epithelial cells. This group includes many of the most common cancers and include nearly all those in the breast, prostate, lung, pancreas and colon.
- **Sarcoma:** Cancers arising from connective tissue (i.e. bone, cartilage, fat, nerve), each of which develops from cells originating in mesenchymal cells outside the bone marrow.
- **Lymphoma and leukemia:** These two classes arise from hematopoietic (blood-forming) cells that leave the marrow and tend to mature in the lymph nodes and blood, respectively.[88]
- **Germ cell tumor:** Cancers derived from pluripotent cells, most often presenting in the testicle or the ovary (seminoma and dysgerminoma, respectively).
- **Blastoma:** Cancers derived from immature "precursor" cells or embryonic tissue.

Cancers are usually named using -carcinoma, -sarcoma or -blastoma as a suffix, with the Latin or Greek word for the organ or tissue of origin as the root. For example, cancers of the liver parenchyma arising from malignant epithelial cells is called hepatocarcinoma, while a malignancy arising from primitive liver precursor cells is called a hepatoblastoma and a cancer arising from fat cells is called a liposarcoma. For some common cancers, the English organ name is used. For example, the most common type of breast cancer is called ductal carcinoma of the breast. Here, the adjective ductal refers to the appearance of cancer under the microscope, which suggests that it has originated in the milk ducts.

Benign tumors (which are not cancers) are named using -oma as a suffix with the organ name as the root. For example, a benign tumor of smooth muscle cells is called a leiomyoma (the common name of this frequently occurring benign tumor in the uterus is fibroid). Confusingly, some types of cancer use the -noma suffix, examples including melanoma and seminoma.

Some types of cancer are named for the size and shape of the cells under a microscope, such as giant cell carcinoma, spindle cell carcinoma and small-cell carcinoma.

CHAPTER II

2. REVIEW OF LITERATURE

Dong-Jiann Huang et al, (2004)¹¹⁹ investigated possible antioxidant and antiproliferative activities of the different extracts from sweet potato (*Ipomoea batatas* [L.] Lam 'Tainong 57') organs. DPPH staining, total phenolic compounds and flavonoid content, DPPH radical, reducing power method, FTC method, and cell proliferation were all employed. In the DPPH staining, ethanol extract of vein had the highest radical-scavenging activity when it was diluted to 6.25 mg dry matter/mL. Among all the extracts, the highest amount of total phenolic and flavonoid compounds was found in the ethanol extract of vein. In the DPPH colorimetric method, it was found that ethanol extract of leaf had the highest radical-scavenging activity, followed by water extract of vein. In the reducing power activity assay, it was found that the water extract of leaf had the highest reducing power activity, followed by ethanol extract of vein. Like phenolic compounds, the highest FTC activity was found in the ethanol extract of vein. The antiproliferative activities of sweet potato were studied in vitro using human lymphoma NB4 cells, and the following results were found: water extract of vein had the highest antiproliferative activity with an EC_{50} of $449.6 \pm 27.73 \mu\text{g/mL}$, followed by water extract of storage root, water extract of leaf, ethanol extract of storage root, and ethanol extract of leaf. Although the ethanol extract of vein showed strong antioxidant activity, it had no antiproliferative activity under the experimental conditions tested.

Prasanth et al (2010)¹²⁰ studied invitro cytotoxic and antioxidant properties of ethanolic extract of *ipomoea batatas*. The extract showed potent cytotoxic activity in trypan blue

e dye exclusion method using DLA cell lines with EC50 value of 305µg/ml and exhibited a dose dependent decrease in cell count for all the concentrations tested. The antioxidant activity was evaluated by DPPH free radical method. The extract exhibited potent antioxidant activity with an EC50 of 36.5µg/ml.

Wamidh H. Talib et al (2010)¹²¹ Reported Forty four extracts from sixteen plants used traditionally as anticancer agents in vitro for their antiproliferative activity against Hep-2, MCF-7, and Vero cell lines. Plants were fractionated using ethanol, methanol, chloroform, n-hexane, distilled water, and butanol. The antiproliferative activity was measured by MTT assay. TLC was used to identify active fractions. The apoptotic activity of active fractions was determined using TUNEL colorimetric assay. 20 of these extracts demonstrated significant antiproliferative activity against one or more of the cell lines. These extracts were prepared from *Ononishirta*, *Inulaviscosa*, *Salvia pinardi*, *Verbascumsinaiticum* and *Ononissicula*. Methanol fractions of *Ononishirta*(aerial parts) and *Inulaviscosa*(flowers) were the most active fractions against MCF-7 cells with IC50 of 27.96 and 15.78 µg/ml respectively and they were less toxic against other cell lines. Other extracts showed lower activity against cancer cell lines. TLC analysis showed the presence of flavonoids and terpenoids in active plants while alkaloids were detected in *Ononishirta*(aerial parts) extracts. *Ononishirta*(aerial parts) and *Inulaviscosa*(flowers) extracts exerted their antiproliferative activity by inducing apoptosis in cancer cell lines.

Seow-Mun Hue et al (2011)¹²² Evaluated *Ipomoea batatas* (Sweetpotato) is currently ranked sixth in the total world food production and are planted mainly for their storage roots. The present study was undertaken to evaluate and compare the antioxidant properties of the leaf and carotenoids extract from

the *Ipomoea batatas* var. Oren leaves. Total flavonoids in the leaf extract was $144.6 \pm 40.5 \mu\text{g/g}$ compared to $114.86 \pm 4.35 \mu\text{g/g}$ catechin equivalent in the carotenoids extract. Total polyphenols in the leaf extracts ($3.470 \pm 0.024 \text{ GAE g/100g DW}$) was slightly higher compared to carotenoids extract ($2.994 \pm 0.078 \text{ GAE g/100g DW}$). The carotenoids extract marked a higher radical scavenging capacity with the $\text{IC}_{50} = 491.86 \mu\text{g/ml}$ compared to leaf extract ($\text{IC}_{50} = 545.39 \mu\text{g/ml}$). Concentration-dependent reducing activity was observed for both extracts. Thus, the carotenoids extraction process retained most of the antioxidant capacity from the leaves and can be made into potential natural yellow dye with antioxidant property.

Vandana Panda et al, (2011)¹²³ Evaluated *Ipomoea batatas* (L.) Lam. from the family Convolvulaceae is the world's sixth largest food crop. The tubers of *Ipomoea batatas* commonly known as sweet potato are consumed as a vegetable globally. The tubers contain high levels of polyphenols such as anthocyanins and phenolic acids and vitamins A, B and C, which impart a potent antioxidant activity that can translate well to show wound healing effects. To check their effects on wound healing, the peels and peel bandage were tested on various injury models in rats in the present study. The methanolic extracts of the peels and peel bandage of *Ipomoea batatas* tubers (sweet potato) were screened for wound healing by excision and incision wound models on Wistar rats. Three types of gel formulations were prepared, viz., gel containing 3.0% (w/w) peel extract, gel containing 6.0% (w/w) peel extract and gel containing 10% (w/w) peel extract. Betadine (5% w/w povidone iodine cream) was used as a reference standard. In the incision wound model, Tensile strength of the skin was measured. Epithelization time, wound contraction, hydroxyproline content of the scab, and ascorbic acid and malondialdehyde content of the plasma were determined in the excision wound model. Results: In the incision

wound model, high tensile strength of the wounded skin was observed in animals treated with the peel extract gels and the peel bandage when compared with wounded control animals. The increase in tensile strength indicates the promotion of collagen fibers and that the disrupted wound surfaces are being firmly knit by collagen. In the excision wound model, significant wound closure was observed on the 4th day in rats treated with all three gel Formulations when compared with the wounded control rats. A significant increase in hydroxyproline and ascorbic acid content in the gel-treated animals and a significant decrease in malondialdehyde content in the animals treated with gel as well as peel bandage was observed when compared with the wounded control animals. It may be concluded that the peels of *Ipomoea batatas* tubers possess a potent wound healing activity, which may be due to an underlying antioxidant mechanism.

Vandana Panda et al, (2012)¹²⁴ Conducted studies on peptic ulcers occur in that part of the gastrointestinal tract which is exposed to gastric acid and pepsin, i.e., the stomach and duodenum. Gastric and duodenal ulcers are common pathologies that may be induced by a variety of factors such as stress, smoking and noxious agents including non-steroidal anti-inflammatory drugs. *Ipomoea batatas* tubers (sweet potato) contain ample amounts of antioxidants. It has been proven already by many scientific studies that antioxidants have ulcer healing properties. In reference to this, we tried assessing the ulcer healing effect of *Ipomoea batatas* tubers. The anti-ulcer activity of the tubers of *Ipomoea batatas* (sweet potato) was studied in cold stress and aspirin-induced gastric ulcers in Wistar rats. Methanolic extracts of *Ipomoea batatas* tubers (TE) at two doses, viz., 400 and 800 mg /kg were evaluated in cold stress and aspirin-induced gastric ulcer models using cimetidine

and omeprazole respectively as standards. The standard drugs and the test drugs were administered orally for 7 days in the cold stress model and for 1 day in the aspirin-induced gastric ulcer model. Gastroprotective potential, status of the antioxidant enzymes {superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPx) and glutathione reductase (GR)} along with GSH, and lipid peroxidation were studied in both models. Results: The results of the present study showed that TE possessed gastroprotective activity as evidenced by its significant inhibition of mean ulcer score and ulcer index and a marked increase in GSH, SOD, CAT, GPx, and GR levels and reduction in lipid peroxidation in a dose dependant manner. Conclusion: The present experimental findings suggest that tubers of *Ipomoea batatas* may be useful for treating peptic ulcers.

Yuzhijiao et al (2012)¹²⁵ The radical scavenging effects by α,α -diphenyl- β -picrylhydrazyl (DPPH) and superoxide anions of anthocyanin extract from purple sweet potato were investigated. The antioxidation experiments showed that the reducing power of the anthocyanin extract reduced 0.572 at 0.5 mg/ml, while those of Lascorbic acid (L-AA) and butylatedhydroxytoluene (BHT) reduced 0.460 and 0.121, respectively. They also displayed potent antioxidant effects against the DPPH radical and superoxide anions radical, showing the IC₅₀ values of 6.94 and 3.68 μ g/ml, respectively. Moreover, this anthocyanin extract also could significantly inhibit the formation of lipid peroxidation compound. Sixteen kinds of anthocyanins in purple sweet potato were detected by high-performance liquid chromatography with diode-array detection (HPLC-DAD), and most of the anthocyanins were acylated.

Marilena Meira et al (2012)¹²⁶ ,Approximately 600-700 species of *Ipomoea*, Convolvulaceae, are found throughout tropical and

subtropical regions of the world. Several of those species have been used as ornamental plants, food, medicines or in religious ritual. The present work reviews the traditional uses, chemistry and biological activities of *Ipomoea* species and illustrates the potential of the genus as a source of therapeutic agents. These species are used in different parts of the world for the treatment of several diseases, such as, diabetes, hypertension, dysentery, constipation, fatigue, arthritis rheumatism, hydrocephaly, meningitis, kidney ailments and inflammations. Some of these species showed antimicrobial, analgesic, spasmolytic, spasmogenic, hypoglycemic, hypotensive, anticoagulant, anti-inflammatory, psychotomimetic and anticancer activities. Alkaloids, phenolics compounds and glycolipids are the most common biologically active constituents from these plant extracts.

Adsull et al (2012)¹²⁷ Microbial diseases remain a major challenge for modern science even today. Natural products are used as traditional medicines from ancient times. They are having a great importance in Ayurveda. One of the medicinal plant species is *Ipomoea carnea* belongs to convolvulaceae family and *fistulosa* sub-family. Antimicrobial activity was tested against n-hexane, ethyl acetate, acetone, ethanol and acetone fraction of acetone extract. The investigation is carried out against various gram positive and gram negative bacterial strains (*Escherichia coli* ATCC – 11246; *Staphylococcus aureus* ATCC – 6538 P; *Salmonella typhimurium* ATCC – 23564; *Pseudomonas aeruginosa* ATCC – 27853; *Proteus vulgaris* ATCC – 13315; *Bacillus cereus* ATCC – 11778) . Disc diffusion method is employed for the detection of antimicrobial activity. Streptomycin was used as standard. The crude acetone extract (3) exhibits activity against *Proteus vulgaris* and *Salmonella typhimurium*, while the crude ethanol extract (4) elucidates antimicrobial activity against *Pseudomonas aeruginosa*

RajuAsirvatham et al (2013)¹²⁸ Conducted in vitro studies was performed to examine the antioxidant and anticancer activities of ethanol and aqueous extracts of *Drosera indica* L. Methods: Different concentrations (5 – 640 mcg/ml) of the ethanol (EEDI) and aqueous (AEDI) extracts of *D. indica* L were used in various antioxidant assay methods such as hydroxyl radicals, DPPH, super oxide radical scavenging activity, chelating ability of ferrous ion, nitric oxide radical inhibition, ABTS and reducing power. Ascorbic acid (AA) was used as the standard antioxidant for the free radical scavenging assays. Dalton's Ascitic Lymphoma (DAL) and Ehrlich Ascitic Carcinoma (EAC) cell lines were used as the in vitro cancer models for the trypan blue dye and LDH leakage assays, where 5 to 250 mcg /ml of both EEDI and AEDI were tested. Results: EEDI showed antioxidant activities with the minimum IC₅₀ values of 34.8±0.43 mcg/ml in scavenging of hydroxyl radical and moreover AEDI showed minimum IC₅₀ values of 94.51±0.84 mcg/ml in Fe²⁺ chelating assay. EEDI on the reducing power assay and ABTS showed higher IC₅₀ than standard AA. IC₅₀ values of AEDI on Fe²⁺ chelating assay and super oxide radical assay was lesser than IC₅₀ value of AA. Both extracts at 250 mcg/ml dose showed remarkable increase in the percentage of dead cancer cells (90% by EEDI and 86% by AEDI in DAL model and 89% by EEDI and 80% by AEDI in EAC model).

Eleazu et al(2013)¹²⁹ Evaluated The physicochemical composition, functional properties, inhibitory actions and energy value of the flour of a cream fleshed sweetpotato variety (TIS/87/0087) that is high yielding and commercially sold in South Eastern Nigeria were investigated using standard techniques. The flour was observed to have good functional properties with a pH of 5.32±0.01, high percentage moisture content, indicative of poor shelf life characteristics and high chances of being attacked by microbes, low percentage dry

matter, lipid, crude fibre and ash contents but a promising source of starch ($20.78 \pm 0.02\%$), carotene ($5.0 \pm 0.04 \mu\text{g/g}$), protein ($2.67 \pm 0.59\%$), carbohydrate ($40.77 \pm 3.05\%$), energy ($179.61 \pm 20.97 \text{ kcal/100 g}$), polyphenols, in addition to containing significant quantities of reducing sugar ($1.58 \pm 0.53\%$). In addition, the methanolic extract of the flour possessed higher scavenging activities on 2,2 diphenyl-1-picrylhydrazyl (DPPH) radical than standard quercetin. Results show that this sweet potato variety has potentials of biological properties and could have wide utility in food, alcohol and sugar industries. In addition, it could serve as a promising source of protein and its consumption could be utilized in the management of diseases that implicate free radicals. Finally, it could also be useful as a drug binder and disintegrant in pharmaceutical industries.

Zainal Baharum et.al (2014)¹³⁰ The aims of this study were to determine the antioxidant and antiproliferative activity of the following *Theobroma cacao* plant part methanolic extracts: leaf, bark, husk, fermented and unfermented shell, pith, root, and cherelle. Antioxidant activity was determined using 2,2-diphenyl-2-picrylhydrazyl (DPPH), thiobarbituric acid-reactive substances (TBARS), and Folin-Ciocalteu assays; diphenyltetrazolium (MTT) assay was used to determine antiproliferative activity. The root extract had the highest antioxidant activity; its median effective dose (EC_{50}) was $358.3 \pm 7.0 \mu\text{g/mL}$ and total phenolic content was $22.0 \pm 1.1 \text{ g GAE/100 g extract}$ as compared to the other methanolic plant part extracts. Only the cherelle extract demonstrated $10.4\% \pm 1.1\%$ inhibition activity in the lipid peroxidation assay. The MTT assay revealed that the leaf extract had the highest antiproliferative activity against MCF-7 cells [median inhibitory concentration (IC_{50}) = $41.4 \pm 3.3 \mu\text{g/mL}$]. Given the overall high IC_{50} for the normal liver cell line WRL-68, this study indicates that *T. cacao* methanolic extracts have a cytotoxic effect in cancer cells, but not in normal cells. Planned

future investigations will involve the purification, identification, determination of the mechanisms of action, and molecular assay of *T. cacao* plant extracts.

Marcelia Sugata et.al (2015)¹³¹ Purple-fleshed sweet potato (PFSP) Tainung 73 possesses high amount of antioxidative compounds, such as phenolics, flavonoid, and anthocyanin. The major anthocyanin is cyanidin or/and peonidin and their acylated derivatives. Study on the possible properties of PFSP extracts showed that these extracts had potential anti-inflammatory and anticancer activities. Anthocyanin-rich extracts of PFSP TNG 73 could suppress the production of nitric oxide (NO) and some proinflammatory cytokines, such as NF κ - β , TNF- α , and IL-6, in LPS-induced macrophage cell. Nevertheless, these extracts showed no cytotoxicity effect on macrophage cells. On the other hand, these extracts could inhibit the growth of some cancer cell lines, such as human breast cancer (MCF-7), gastric cancer (SNU-1), and colon adenocarcinoma (WiDr), in concentration- and time-dependent manner. After further investigation on molecular mechanism, PFSP TNG 73 extracts demonstrated the ability to induce apoptosis in MFC-7 cancer cell line through extrinsic and intrinsic pathways. Thus, PFSP TNG 73 can be used for future application of drugs, nutritional food, and health supplement.

Milind parle et al (2015)¹³² Sweet potato is an extremely versatile and delicious vegetable that posses high nutritional value. Sweet potato has been grown in tropical and subtropical regions throughout the world, since ancient times. From the times immemorial, the whole sweet potato plant including leaves, stem, and tuberous root is used as traditional medicine. Nowadays, Sweet potato is preferred over other vegetables due to its multifaceted medicinal properties. The medicinal properties of sweet potato include anti-cancer, anti-inflammatory, anti-

diabetic, anti-oxidant, anti-bacterial, anti-fungal, anti-viral, anti-ulcer, hepatoprotective, wound healing and immunomodulatory activities. Sweet potatoes contain magnesium, a crucial mineral, which promotes relaxation, calmness and nerve health. Overall objective of this review article is to give a brief knowledge about the nutritional value, health benefits, phytochemical composition, pharmacological actions and medicinal properties of sweet potato. Sweet potato holds first rank (super food) in nutrition among vegetables.

HuaJi et al (2015)¹³³ In this study, we selected four different color fleshed sweet potatoes, purple- (Jizi 01), red- (Xinong 431), yellow- (Beijing 553) and white- (Shangshu 19) fleshed cultivars as test materials, analyzed nutrient composition, dietary fiber content, anthocyanins quantification, and total phenolics content, and also measured their total antioxidant activity in four different types of sweet potato. In view of differences in flesh color, the nutrient contents of different cultivars appeared to be significantly different. Starch contents of Beijing 553 and Shangshu 19 were higher, but fat contents were lower than others. Protein content of Shangshu 19 was the highest followed by Jizi 01 and Xinong 431. In addition, our analysis results confirmed that purple fleshed sweet potato possesses much higher anthocyanins content than others, even up to 6.23 mg/g dry matter. Also, dietary fiber, total phenolics content, and total antioxidant capacity of Jizi 01 were significantly higher.

Sucharitha et al (2016)¹³⁴ A diuretic is any substance that promotes the production of urine. This includes forced diuresis. There are several categories of diuretics. All diuretics increase the excretion of water from bodies, although each class does so in a distinct way. Alternatively, an ant diuretic such as vasopressin, or ant diuretic hormone, is an agent or drug which

reduces the excretion of water in urine. In medicine, diuretics are used to treat heart failure, liver cirrhosis, hypertension, influenza, water poisoning, and certain kidney diseases. Some diuretics, such as acetazolamide, help to make the urine more alkaline and are helpful in increasing excretion of substances such as aspirin in cases of overdose or poisoning. Diuretics are often abused by those with eating disorders, especially bulimics, in attempts to lose weight. The antihypertensive actions of some diuretics (thiazides and loop diuretics in particular) are independent of their diuretic effect. That is, the reduction in blood pressure is not due to decreased blood volume resulting from increased urine production, but occurs through other mechanisms and at lower doses than that required to produce diuresis. Indapamide was specifically designed with this in mind, and has a larger therapeutic window for hypertension (without pronounced diuresis) than most other diuretics. The main objective of the present research work is to isolate the bioactive molecules and evaluate the diuretic activity of aqueous extract of Ipomoea batata. The phytochemical analysis of aqueous extract of Ipomoea batata root showed the presence of various phytochemical constituents such as flavonoids, carbohydrates, tannins, phenol. The effect of aqueous extract of root of Ipomoea batata on rats with reference to biochemical changes in serum. The group-II (Standard Hydrochlorothiazide 10 ml/kg, p. o) animals showed significant ($P < 0.01$) increase in total urine volume ml/100 gm/hr (10.44 ml). Whereas animals received AEIB significantly ($P < 0.01$) increase in total urine volume ml/100 gm/hr (4.44 and 8.06 ml) and significantly ($P < 0.05$) increased total 200 & 400 mg/kg doses respectively. The phytochemical studies revealed the presence of Carbohydrate, flavonoids, Tannins in the AEIB these may be responsible for its pharmacological activities.

CHAPTER III

.3. AIM AND OBJECTIVE

Cancer is one of the major problem in worldwide due to life style, usage of cancer medications, tobacco usage, lack of proper medication, and undefined drug targets. The treatment of cancer after so many years of research and experience is still unsatisfactory due to certain special characteristics of the cancer cell like capacity for uncontrolled proliferation, invasiveness, metastasis and dedifferentiation. Such dedifferentiated cancer cells can multiply faster when compared to well differentiated cancer cells. Several modern drugs are available in the market do not fulfill the requirements and also with many side effects. So the priority is drugs which minimize the side effects and also required chemical entity for the treatment of cancer with specific action. Several literatures indicated traditional herbs possessing lesser side effects compared to synthetic drugs. The herbal formulations which developed from ayurveda, traditional system of Indian medicine and its additional system of medicine, has been found to have anticancer activity. *Ipomoea batatus* is a traditionally used herb which posses anticancer activity but no scientific validation.. Some scientific studies show that leaf and root extract of *ipomoea batatus* posses anticancer activity. But there was no literature review about anticancer property of the stem available. So the present study is an attempt to develop plant based anticancer drug which will be lesser side effect.

CHAPTER IV

4. PLAN OF WORK

- I. literature Surevey
- II. Extraction of dried stem of the *Ipomoea batatus* by using Soxhlet Apparatus.
- III. Preliminary phytochemical screening of the ethanolic extract of the plant using specific chemical test.
- IV. Perform *invitro* Anti cancer activity will carried out by effect of ethanolic extract of the stem of *ipomoea batatus* on MTT assay.
- V. Selection of fraction with higher activity.
- VI. The *Invivo* anticancer activity of ethanolic extract of stem of *Ipomoea batatus* by using DLA Method in mice.
- VII. Estimating the effect of ethanolic extract of stem of *ipomoea batatus* on
 - Mean survival time
 - Hemoglobin Count
 - RBC Count
 - WBC Count
- VIII. Confirming the anticancer activity of active constituents from literarture.
- IX. Conclusion of the work.

CHAPTER V
5. PLANT PROFILE



5.1. Plant profile

Kingdom : *plantae*

Subkingdom : *Tracheobionta*

Super division: *Spermatophyte*

Division : *Sagnoliophyta*

Class : *Magnoliopsida*

Sub class : *Asteridae*

Order : *Solanales*

Family : *Convolvulaceae*

Genus : *Ipomoea* L.

Species : *I. batatas* (L.) LAM

Synonyms :

Kannada : *Genasu*.

Hindi : *Shakarkand / Ratalu*.

Telugu : *Chilakada dumpa*.

Marathi : *Ratala*.

Bengali : *MishtiAlu*.

Malayalam : *Mathura Kizhangu*.

Scientific names

batatasedulis Choisy (Kamote (all dialects))

Convolvulus batatas Linn (Lapin (if.))

Convolvulus edulis Choisy (Pangg-bagun (Sul.))

Ipomoea batatas (L.) (Sweet potato (Engl.),

(Yam (Engl.),

5.2. Geographical distribution:

The plant is commonly seen growing in all parts of India, Asian and some other European and American countries(9).

5.3. Chemical constituents of leaf:

Sweetpotato roots and tops possess a variety of chemical compounds relevant to human health. About 80 to 90 % of sweetpotato dry matter is made up of carbohydrates, consisting mainly of starch (60-70%) and sugars with lesser amounts of pectins, hemicelluloses and cellulose. Sweet-potato also contains

protein (0.46%-2.93%), dietary fiber (0.49%-4.71%), lipid (0.06%-0.48%) and ash (0.31%-1.06%).

It contains essential mineral nutrients such as Ca(117mg), (0.56mg), vit E(0.56mg), Fe(1.8mg)/100mg, S, Cu, Zn, P, Mg, Na, K, Mn, Al and B. Sweetpotato is also an important source of vitamin A, thiamin, riboflavin, niacin, ascorbic acid, β -carotene and many other functional compounds.

Sweetpotato leaves are an excellent source flavonoids with antioxidative poly-phenols, with 6 polyphenolic compounds, 15 anthocyanins and phenolic acids such as caffeic, moncaffeoylquinic (chloro-genic), dicaffeoylquinic and tricaffeoylquinic acids, and are superior in this regard to other commercial vegetables.

The another major constituent flavonoids is proanthocyanins and two or more flavan-3-ol such as catechin, epicatechin or gallocatechin. Catechin contains 2 benzene rings to be the powerful scavenger(8).

5.4. Uses Of Ipomoea Batatas

1 :The young leaves and shoots are sometimes eaten as greens.

2 : All parts of the plant are used for animal fodder.

3 : Industrial alcohol production.

4: Sweetpotato leaves used as a vegetable, a tea, in noodles, in breads, in confectioneries and as a nutritional supplement.

Reported activities of Ipomoea batatas:

1: Makoto yoshimoto and shoji yahara evaluated the anti-mutagenic activity of the caffeoylquinic acid derivatives in Ipomoea batatas leaf.

2: poly phenolics have attracted special attention due to their use in oxidative stress which may cause cancer, aging.

3: Mukesh Nandave, SK Ojha and DS Arya reported that flavonoids show anti-thrombotic, anti-ischemic, anti-arrythmic and cardioprotective activity by free radical scavenging mechanism.

4: The leaf extracts also shows anti-bacterial activity, ultraviolet protection effect, anti-inflammation and promotion in bowel movement.

Folkloricuse:

-Crushed leaves applied to boils and acne.

-Boiled roots used for diarrhea.

"Ipomoea" comes from the Greek words ipos, which means "bind weed" and homoios which means "resembling". When this is put together to form "Ipomoea" the direct translation is "resembling bindweed". This name makes sense because the sweet potato has a twining habit, much like the bindweed. The species name "batatas" was originally the Taino name for sweet potato.

Ipomoea batatas is a perennial climber growing up to 3 meters at a fast rate. This is one of those rare plants that are grown as a tasty vegetable and a pretty ornamental. Sweet potato vine is a vigorous, herbaceous, trailing vine, is native to Central America and the Pacific Islands. It offers sweet edible tubers and leaves and attractive foliage.

The leaves are heart-shaped or deeply lobed the colors vary from chartreuse, to bronze or purple and can be variegated. The funnel-shaped flowers can be pale rosy purple or white. They appear when days are long and growing conditions favorable.

CHAPTER VI

6. MATERIALS AND METHODS

6.1. Collection of plant

The stem of Ipomoea batatus were obtained from Department of Horticulture, Malappuram and it was authenticated by Dr.Raghu.A.V Kerala Forest Research Institute Peechi, Trissur.

6.2. Reagents and Requirements for preliminary extraction procedure

1. Reflux apparatus.
- 2.Methanol (Merk Germany)
- 3.Dried stem of Ipomoea batatus
- 4.Glass ware apparatus
- 5.Dessicator

6.3. Extraction of plant material

6.3.1. Extraction 1

About 100 g of powdered sample was weighed and extracted with 500 ml of methanolby reflux method at 24 hours. After the reflux, extract was collected dried under reduced pressure, which was stored in desiccators. Complete moisture was removed from the extract and yield was calculated by using the following formula

$$\text{Percentage yield} = (B/A) \times 100$$

Where

A = Weight of powder

B = Weight of extract

6.3.2. Extraction 2

About 100 g of dried stem was taken, powdered by a grinder, weighed and extracted with 500ml of chloroform by reflux method at 24 hours. After the reflux, extract was collected dried under reduced pressure which was stored in desiccators in order to remove complete moisture from the extract and yield was calculated by using the above formula.

6.3.3. Extraction 3

About 100g of powdered sample was weighed and extracted with 500ml of ethanol by reflux method at 24 hours. After the reflux, the extract was collected dried under reduced pressure, which was stored in desiccators until complete moisture was removed from the extract and yield was calculated by using the above formula.

6.4. Preliminary phytochemical screening of the plant

The ethanol extract of the plant was used for testing preliminary phytochemical screening in order to detect major groups.

6.4.1. Test for carbohydrates

- a. Molich's test:** Dissolve small quantity of ethanolic extract of Ipomoea batatas separately in a 4ml distilled water and filtered. The filtrate was subjected to Molisch's test.
- b. Fehling's test:** Dissolve a small portion of extract in water and treat with Fehling's solution.
- c. Phenols test:** The extract was spotted on a filter paper. A drop of phosphomolybdic acid reagent was added to the spot and was exposed to ammonia vapour.

6.4.2. Test for flavonoids

- a. **Shinoda test:** To the 2 to 3 ml of extract, a piece of magnesium ribbon and 1 ml of concentrated HCL was added.
- b. **Lead acetate test:** To the 5 ml extract 1 ml of lead acetate solution was added.

6.4.3. Test for tannins

- a. **Braemer's test:** to the 2 to 2 ml extract, 10% alcoholic ferric chloride solution was added.

6.4.4. Test for steroid/terpinoid

- a. **Libbermann-Burchardt test:** To 1 ml of extract, 1 ml of chloroform, 2 to 3 ml of acetic anhydride and 1 to 2 drops of concentrated sulphuric acid are added.

6.4.5. Test for alkaloids

- a. **Draggendorfs test:** A drop of extract was spotted on small pieces of precoated TLC plate and the plate was sprayed with modified Draggendorfs reagent
- b. **Hager's test:** The extract was treated with few ml of Hager's reagent.
- c. **Wagner's test:** The extract was treated with few ml of Wagner's reagent.

6.4.6. Test for Glycosides

- a. **Legal's test:** Dissolve the extract in 2 ml pyridine, add sodium nitroprusside solution 2 ml, and made alkaline with sodium hydroxide solution.

6.4.7. Test for Saponins

- a. **Foam test:** 1 ml of extract was diluted with 20ml of distilled water and shaken with a graduated cylinder for 15 minutes.

6.4.8. Test for Amino acids

- a. **Ninhydrin test:** Dissolve a small quantity of extract in few ml of water and added 1ml of ninhydrin reagent.

6.4.9. Test for anthraquinones

- a. **Borntrager's test:** About 50mg of powder extract was treated with 10% of ferric chloride solution and 1 ml of concentrated HCL. The extract was cooled, filtered, and the filtrate was shaken with diethyl ether. The ether extract was further extracted with strong ammonia.

6.4.10. Test for fixed oils and fats

- a. Press small quantity of the petroleum ether extract between two filter paper.

The results of the above experiments can be noted as follows.

- ❖ If the response to the test is high it can be noted as +++ which indicates that the particular group is present as the major class.
- ❖ If the response to the test is average it can be noted as ++ which indicates the presence in moderate quantity.
- ❖ If the response is very small note it as + indicating the presence of only in traces the response is very small
- ❖ If no response is then negative.

6.5. Pharmacological Screening

6.5.1. Invitro Anticancer Activity

6.5.2. Mtt Assay: Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. J Immunol Methods 16;65(1-2):55-63

6.5.3. Tumour Cells

Tumor cells Ehrlich Ascites Carcinoma (EAC) cells were obtained from Amala cancer research center, Trissur, Kerala, India. The cells were maintained in vivo in Swiss albino mice by intraperitoneal transplantation. EAC cells aspirated from the peritoneal cavity of mice were washed with saline and given intraperitoneally to develop ascetic tumor.

6.5.2. Mtt Assay (Helacell Lines)

1. For adherent cells, remove the medium and replace it with 100 μ L of fresh culture medium. For non-adherent cells, centrifuge the microplate, pellet the cells, carefully remove as much medium as possible and replace it with 100 μ L of fresh medium.
2. Add 10 μ L of the 12 mM MTT stock solution (prepared in step 1.1) to each well. Include a negative control of 10 μ L of the MTT stock solution added to 100 μ L of medium alone.
3. Incubate at 37°C for 4 hours. At high cell densities (>100,000 cells per well) the incubation time can be shortened to 2 hours.
4. Add 100 μ L of the SDS-HCl solution (prepared in step 1.2) to each well and mix thoroughly using the pipette.

5. Incubate the microplate at 37°C for 4– hours in a humidified chamber. Longer incubations will decrease the sensitivity of the assay.
6. Mix each sample again using a pipette and read absorbance at 570 nm.

6.5.4. Animals

Male Swiss albino mice weighing 25- 30g were used in the present study. All rats were kept at room temperature of 22-25°C in the animal house. All the animals were followed the internationally accepted ethical guidelines for the care of laboratory animals. Prior to the experiments, rats were fed with standard food for 1 week in order to adapt to the laboratory conditions. All animal procedures were performed after approval from the institutional ethics committee. The experimental protocol has been approved by institutional animal ethics committee, The Karpagam College of Pharmacy, Coimbatore

6.5.5. Acute toxicity study

Ethanollic extract of Ipomoea batatus stem were studied for acute toxicity at doses of 5mg/kg, 50mg/kg, 300mg/kg, 500mg/kg and 2000mg/kg. As per OECD 420 guideline dose of 2000mg/kg showed the toxic symptoms, so according to OECD guideline 420, it is considered as a LD50 cutoff value. Doses selected for pharmacological studies by fixed dose methods are 250mg/kg and 500mg/kg⁸.

6.6. In vivo Anticancer Activity

The Swiss albino mice (20-25 g) were divided into five groups (n=12). Except, Group-I all the animals in DLA group were being injected with DLA cells (2×10^6 cells/mouse, i.p). This was marked as day "0". Group-I was served as normal DLA control and group-II was served as standard group. After 24h, DLA transplanted group-III, IV and V were being injected (100, 200 and 400mg/kg b.w. i.p.) once daily for 12 consecutive days. Group II received standard drug 5-Fluorouracil (20 mg/kg i.p) for 12 consecutive days. After administrations of last dose 6 mice from each group were kept fasting for 18h and blood was collected by direct cardiac puncture for the estimation of haematological determination. Rest of animals in each groups were kept alive with food and water ad libitum to check the percentage increase in life span of the tumor host and also to determine the mean survival time (MST). Antitumor activity of MeIB extract was assessed by observation of changes with respect to the following parameters

6.6.1. Antitumour Parameters

Percentage increase life span The effect of extract on tumor growth was monitored by recording the mortality daily for a period of 6 weeks and percentage increase in average life span was calculated. $\% \text{ ILS} = \{(\text{life span of treated group} / \text{life span of controlled group}) - 1\} \times 100$

6.6.2. Body weight analysis

Body weights of the experimental mice were recorded both in the treated and control groups at the beginning of the experiment (day 0) and sequentially on every 5th day during the treatment period and calculated on 15th day.

6.6.3. Changes in food intake

Feed consumed by 6 animal/cage/week = Total quantity of feed offered during that week (gm) – Feed left over on last day of week (gm). Feed consumed by individual animal/week = Feed consumed by 6 animal per cage per week/6. Feed consumed by individual animal/day = Feed consumed by individual animal per week/7.

6.6.4. Determination of tumor volume and weight

The mice were dissected and the ascetic fluid was collected from the peritoneal cavity. The volume was measured by taking it in a graduated centrifuge tube and weight immediately.

6.6.5. Determination of Hematological parameters

Blood was drawn from each mouse by the retro orbital plexus method and the white blood cells (WBC), red blood cells (RBC), hemoglobin and protein were determined. Serum preparation of the sample blood use to evaluated the biochemical parameters.

6.7. Statistical analysis

All values were expressed as mean \pm SEM. Statistical analysis was performed with one way analysis of variance (ANOVA) followed by Tukey Kramer multiple comparison tests. P value < 0.001 was considered as highly significant and < 0.05 were considered significant when compared to control.

CHAPTER VII

7. RESULTS AND DISCUSSION

7.1 Soxhlet Extraction of *Ipomoea Batatus* Stem

The percentage yield of various extracts viz. petroleum methanol, chloroform and ethanol was 23.98%, 35.83%, 20.46%w/w respectively

Table No:1-The percentage yield of stem of *Ipomoea batatus* stem extracts

Plant	Part used	Method of Extraction	Solvents	Percentage Yield (%W/W)
Ipomoea batatus	Dried stem	Continuous Hot percolation by Soxhlet apparatus	Chloroform	23.98%
			Ethanol	35.83%
			Methanol	20.46%

Figure No: 3 Soxhlet Apparatus



7.2 Preliminary phytochemical screening of *Ipomoea batatus* stem extract

Table No:2 Preliminary phytochemical screening of *Ipomoea batatus* stem extract

Class of compounds	Tests performed	Results
Carbohydrates	Molisch's test Fehling's test	Negative
Phenols	Phosphomolybdic acid test	Positive
Flavonoids	Shinoda test Lead acetate test	Positive
Tannins	Braemer's test	Negative
Alkaloids	Wagner's Mayer's Draggendorf's test	Positive
Glycosides	Legal's test Brontranger's test	Positive
Saponins	Foam test	Positive
Sterols	Salkowski's test	Negative
Aminoacids	Ninhydrin test	Negative
Terpenoids	LibermannBurchardt test	Negative


7.3 *Invitro* Anticancer Activity

The percentage cell viability is reduced with increase in dose of the plant extract.

Table No: 3 Concentration vs Percentage viability

Sample Concentration (µg/ml)	MTT assay –HeLa % viability
Control	96.58±0.25
50	85.26±0.21
100	62.54±0.34
150	56.23±0.38
200	45.23±0.36
250	42.25±0.40

Figure No.3 Effect *Ipomoea batatus* stem extract on percentage cell viability



7.4 Acute Oral Toxicity Studies (OECD 2001)

After administrating the drug we checked the animals for 15 days. The parameters checked includes Mortality rate, Mucous membrane, salivation hair falling etc. all the parameters were found to be normal.

Table No.4 - Common Parameters

Parameters	DLA controll	Standard (cyclophos phamide)	Extract 100mg/k g	Extract 200mg/ kg	Extract 400mg/ kg
Skin and fur	Normal	Normal	Normal	Normal	Normal
Eyes	Normal	Normal	Normal	Normal	Normal
Mucous membrane	Normal	Normal	Normal	Normal	Normal
Behavioural patterns	Normal	Normal	Normal	Normal	Rapid heart beat
COMA	Normal	Normal	Normal	Normal	Normal

Mortality rate is found to be zero percentage. None of the animals died in between the experiment. It shows the extract posses lesser side effect and can be used safely.

Table No: 5 - Mortality rate

Group	Dose (mg/kg)	No.of animals	Dose difference (a)	Mortality Rate (b)
1.	5	6	0	Zero Percentage mortality rate
2.	50	6	45	
3.	300	6	250	
4.	2000	6	1750	

$LD_{50} = \text{higher dose} - \Sigma (a \times b) / n$ Where $n = \text{No. of animals in each group}$

$LD_{50} = 2000 - 0 = 2000 \text{ mg / kg}$


$ED_{50} = LD_{50} / 10 = 2000 / 10 = 200 \text{mg / kg}$

7.5 *Invivo* Anticancer Studies

Table No:7 Effect Of Ipomoea Batatus Stem Extract On Mean Survival Time

PARAM ETERS	CONTROL	STANDARD (20mg/kg)	EXTRACT (100mg/kg)	EXTRACT (200mg/kg)	EXTRACT (400mg/kg)
Mean survival time (days)	14.5±0.14	14±0.14***	13.8±0.28*	13.1±0.42*	12.9±0.28***

Figure No: 7- Effect Of Ipomoea Batatus Stem Extract On Mean Survival Time



7.6 Effect Of *Ipomoea Batatus* Stem Extract On Hemoglobin


The *invivo* immunomodulatory activity results have been expressed that the ethanolic extract of stem of *ipomoea batatus* shown anticancer activity action when compared to control and positive control which is used in dose levels at 100, 200 and 400mg/kg have dose dependend anti cancer activity due to constantly decreased Hemoglobin. From the figure the anti cancer activity conducted on ethanolic extract of fruits of *ipomoea batatus* on Hemoglobin count in mice showed the different drug concentration is less activity as compared to standard drug. After 15 days of analysis the Hemoglobin count is less in different drug concentration as compared to control drug.

Table No.8 Effect Of *Ipomoea Batatus* Stem Extract On Hemoglobin

Group N=6	Treatment	Dose (mg/kg body weight)	Hemoglobin count		
			Day1	Day 10	Day 15
Group I	controll	0.6%	9.0±0.13	7.5±0.15	6.6±0.16
Group II	cyclophosphamide	30	8.9±.018	7.0±0.19	6.1±0.17
Group III	Ethanolic extract	100	8.2±0.22	6.9±0.21	6.0±0.22
Group IV	Ethanolic extract	200	7.9±0.26	6.4±0.25	5.9±0.26
Group V	Ethanolic extract	400	8.5±0.29	7.2±0.30	6.0±0.31

All values are expressed as mean ± SEM evaluated by ANOVA followed by Dunnet's t test

Figure No: 9 Effect Of Ipomoea Batatus Stem Extract On Hemoglobin Count



Drug concentration vs Hb concentration

7.7 Effect Of *Ipomoea Batatus* Stem Extract On RBC


The *invivo* anticancer activity results have been expressed that the ethanolic extract of stem of *ipomoea batatus* shown anticancer activity, when compared to control and positive control which used dose levels at 100,200 and 400mg/kg have dose dependent anticancer activity due to constantly decreased RBC cells

Table No:10 Effect Of *Ipomoea Batatus* Stem Extract On RBC Count

Group (N=6)	Treatment	Dose (mg/kg Body weight)	RBC COUNT (x 10 ⁶ /μL)		
			Day 1	Day 10	Day 15
Group I	Control	6%	7.5±0.32*	6.3±0.35	6.0±0.30
Group II	Cyclophosphamide	20	7.2±0.36*	6.1±0.34	5.8±0.29
Group III	Ethanolic extract	100	7.5±0.40*	6.2±0.38***	5.6±0.35
Group IV	Ethanolic extract	200	7.4±0.38*	6.1±0.33*	5.5±0.28
Group V	Ethanolic extract	400	7.3±0.36*	6.0±0.32**	5.8±0.30

All values are expressed as mean ± SEM evaluated by ANOVA followed by Dunnet's t test

Figure No:10 Effect Of *Ipomoea Batatus* Stem Extract On RBC Count



7.8 Effect Of *Ipomoea Batatus* Stem Extract On WBC


The invivo anticancer activity results have been expressed that the ethanolic extract of stem of ipomoea batatus shown anticancer activity, when compared to control and positive control which used dose levels at 100,200 and 400mg/kg have dose dependent anticancer activity due to constantly decreased WBC cells

Table No:11 Effect Of *Ipomoea Batatus* Stem Extract On WBC

Group N=6	Treatment	Dose (mg/kg body weight)	Total WBC count (cells/mm ³)		
			Day 1	Day 10	Day 15
Group I	Control	0.6%	10442±8.33	10400±0.004	10343±6.6
Group II	Cyclophosphamide	30	10900±0.001*	9900±0.006*	5440±9.6*
Group III	Ethanolic extract	100	9700±0.007*	7250±0.003*	4265±1.00*
Group IV	Ethanolic extract	200	9883±16.67*	7183±0.65*	4060±1.00*
Group V	Ethanolic extract	400	10033±21.08*	6100±0.004*	3900±0.009*

All values are expressed as mean ± SEM evaluated by ANOVA followed by Dunnet's t test

Figure No: 11 Effect Of Ipomoea Batatus Stem Extract On WBC



Drug concentration vs WBC count

CHAPTER VIII

8.SUMMARY AND CONCLUSION

The preliminary phytochemical investigation revealed the compounds present in the stem of *Ipomoea batatus*. *Invitro* cancer studies were performed using MTT assay with cell lines and found that the material possess significant activity. The LD 50 values were also determined by doing the assay in various concentrations.

Herb based drug industry is growing day by day. The destruction of natural habitats over exploitation and unsustainable harvest has lead to a severe scarcity of raw material. These problems have adversely affected the quantity of herbal drugs.

The current work was specific in the field of oncology and revealed the fact that the medicinal plants are capable of treating cancer

The current work revealed that the stem shows significant anti-cancer activity. So the production of a herbal formulation for the disease will be a boon for the entire pharma industry. The identification of chemical compounds present in the stem of *Ipomoea batatus* which are responsible for the specific activity is remaining. In future work I would like to separate the main constituents that present in the plant, responsible for the action and structure elucidation of those constituents. The present study is an attempt to develop novel plant based anti cancer drugs.

CHAPTER IX

9.BIBLIOGRAPHY

1. "Cancer Fact sheet N°297". 2014 World Health Organization. February 2014.
2. "Defining Cancer". 2014 National Cancer Institute. Retrieved 10 June 2014.
3. "Cancer - Signs and symptoms". 2014 NHS Choices. Retrieved 10 June 2014.
4. "Obesity and Cancer Risk".2012 National Cancer Institute. January 3, 2012.
5. Anand P, Kunnumakkara AB, Kunnumakara AB, Sundaram C, Harikumar KB, Tharakan ST, Lai OS, Sung B, Aggarwal BB September 2008. "Cancer is a preventable disease that requires major lifestyle changes". *Pharmaceutical Research*. 25 :2097–116. doi:10.1007/s11095-008-9661-9. PMC 2515569 PMID 18626751.
6. World Cancer Report 2014. World Health Organization.. pp. Chapter 1.1. ISBN 9283204298.
7. "Heredity and Cancer". 2013 American Cancer Society. Retrieved July 22, 2013.
8. "How is cancer diagnosed?". American Cancer Society. 2013-01-29. Retrieved 10 June 2014.
9. Kushi LH, Doyle C, McCullough M, Rock CL, Demark-Wahnefried W, Bandera EV, Gapstur S, Patel AV, Andrews K, Gansler T 2012. "American Cancer Society Guidelines on nutrition and physical activity for cancer prevention: reducing the risk of cancer with healthy food choices and physical activity". *Ca*. 62 1: 30–67. doi:10.3322/caac.20140. PMID 22237782.
10. World Cancer Report 2014. World Health Organization. 2014. pp. Chapter 4.7. ISBN 9283204298.
11. Gøtzsche PC, Jorgensen KJ June 2013. "Screening for breast cancer with mammography". *The Cochrane Database*

- SystematicReviews.6.6:CD001877. doi:10.1002/14651858.CD001877.pub5. PMID 23737396.
12. "Targeted Cancer Therapies". NCI. 2014-04-25. Retrieved 11 June 2014.
 13. World Cancer Report 2014. World Health Organization. 2014. pp. Chapter 1.3. ISBN 9283204298.
 14. "SEER Stat Fact Sheets: 2014 All Cancer Sites". National Cancer Institute. Retrieved 18 June 2014.
 15. "The top 10 causes of death Fact sheet N°310". WHO. May 2014. Retrieved 10 June 2014.
 16. Dubas LE, Ingraffea A February 2013. "Nonmelanoma skin cancer". *Facial Plastic Surgery Clinics of North America*. 21. 1: 43–53. doi:10.1016/j.fsc.2012.10.003. PMID 23369588.
 17. Cakir B, Adamson P, Cingi C November 2012. "Epidemiology and economic burden of nonmelanoma skin cancer". *Facial Plastic Surgery Clinics of North America*. 20 (4): 419–22. doi:10.1016/j.fsc.2012.07.004. PMID 23084294.
 18. Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D (February 2011). "Global cancer statistics". *Ca*. 61 (2): 69–90. doi:10.3322/caac.20107. PMID 21296855.
 19. World Cancer Report 2014. World Health Organization. 2014. pp. Chapter 6.7. ISBN 9283204298.
 20. "Cancer Glossary". cancer.org. American Cancer Society. Retrieved September 11, 2013.
 21. "What is cancer?". cancer.gov. National Cancer Institute. Retrieved September 11, 2013.
 22. Hanahan D, Weinberg RA (January 2000). "The hallmarks of cancer". *Cell*. 100 (1): 57–70. doi:10.1016/S0092-8674(00)81683-9. PMID 10647931.
 23. Hanahan D, Weinberg RA (January 2000). "The hallmarks of cancer". *Cell*. 100 (1): 57–70. doi:10.1016/S0092-8674(00)81683-9. PMID 10647931.

24. Hanahan D, Weinberg RA (March 2011). "Hallmarks of cancer: the next generation". *Cell*. 144 (5):646-74. doi:10.1016/j.cell.2011.02.013. PMID 21376230.
25. Holland Chp. 1
26. Anguiano L, Mayer DK, Piven ML, Rosenstein D (Jul–Aug 2012). "A literature review of suicide in cancer patients". *Cancer Nursing*. 35 (4):E1426. doi:10.1097/NCC.0b013e31822fc76c. PMID 21946906.
27. O'Dell, edited by Michael D. Stubblefield, Michael W. (2009). *Cancer rehabilitation principles and practice*. New York: Demos Medical. p. 983. ISBN 978-1-933864-33-4.
28. Kravchenko J, Akushevich I, Manton KG (2009). *Cancer mortality and morbidity patterns in the U. S. population: an interdisciplinary approach*. Berlin: Springer. ISBN 0-387-78192-7. The term environment refers not only to air, water, and soil but also to substances and conditions at home and at the workplace, including diet, smoking, alcohol, drugs, exposure to chemicals, sunlight, ionizing radiation, electromagnetic fields, infectious agents, etc. Lifestyle, economic and behavioral factors are all aspects of our environment.
29. Tolar J, Neglia JP (June 2003). "Transplacental and other routes of cancer transmission between individuals". *Journal of Pediatric Hematology/Oncology*. 25 (6): 430–4. doi:10.1097/00043426-200306000-00002. PMID 12794519.
30. Biesalski HK, Bueno de Mesquita B, Chesson A, Chytil F, Grimble R, Hermus RJ, Köhrle J, Lotan R, Norpoth K, Pastorino U, Thurnham D (1998). "European Consensus Statement on Lung Cancer: risk factors and prevention. Lung Cancer Panel". *Ca*. 48 (3): 167–76; discussion 164–6. doi:10.3322/canjclin.48.3.167. PMID 9594919.

31. Kuper H, Boffetta P, Adami HO (September 2002). "Tobacco use and cancer causation: association by tumour type". *Journal of Internal Medicine*. 252 (3): 206–24. doi:10.1046/j.1365-2796.2002.01022.x. PMID 12270001.
32. Kuper H, Adami HO, Boffetta P (June 2002). "Tobacco use, cancer causation and public health impact". *Journal of Internal Medicine*. 251 (6): 455–66. doi:10.1046/j.1365-2796.2002.00993.x. PMID 12028500.
33. Sasco AJ, Secretan MB, Straif K (August 2004). "Tobacco smoking and cancer: a brief review of recent epidemiological evidence". *Lung Cancer*. 45 Suppl 2: S3–9. doi:10.1016/j.lungcan.2004.07.998. PMID 15552776.
34. Thun MJ, Jemal A (October 2006). "How much of the decrease in cancer death rates in the United States is attributable to reductions in tobacco smoking?". *Tobacco Control*. 15 (5): 345–7. doi:10.1136/tc.2006.017749. PMC 2563648 . PMID 16998161.
35. Dubey S, Powell CA (May 2008). "Update in lung cancer 2007". *American Journal of Respiratory and Critical Care Medicine*. 177 (9): 941–6. doi:10.1164/rccm.200801-107UP. PMC 2720127 . PMID 18434333.
36. Schütze M, Boeing H, Pischon T, Rehm J, Kehoe T, Gmel G, Olsen A, Tjønneland AM, Dahm CC, Overvad K, Clavel-Chapelon F, Boutron-Ruault MC, Trichopoulou A, Benetou V, Zylis D, Kaaks R, Rohrmann S, Palli D, Berrino F, Tumino R, Vineis P, Rodríguez L, Agudo A, Sánchez MJ, Dorronsoro M, Chirlaque MD, Barricarte A, Peeters PH, van Gils CH, Khaw KT, Wareham N, Allen NE, Key TJ, Boffetta P, Slimani N, Jenab M, Romaguera D, Wark PA, Riboli E, Bergmann MM (April 2011). "Alcohol attributable burden of incidence of cancer in eight European countries based on results from prospective cohort study". *Bmj*. 342:




d1584. doi:10.1136/bmj.d1584. PMC 3072472
. PMID 21474525.





37. Irigaray P, Newby JA, Clapp R, Hardell L, Howard V, Montagnier L, Epstein S, Belpomme D (December 2007). "Lifestyle-related factors and environmental agents causing cancer: an overview". *Biomedicine & Pharmacotherapy = Biomedecine & Pharmacotherapie*. 61 (10): 640–58. doi:10.1016/j.biopha.2007.10.006. PMID 18055160.
38. "WHO calls for prevention of cancer through healthy workplaces"(Press release).2007 World Health Organization. 27 April 2007.
39. Kushi LH, Byers T, Doyle C, Bandera EV, McCullough M, McTiernan A, Gansler T, Andrews KS, Thun MJ (2006). "American Cancer Society Guidelines on Nutrition and Physical Activity for cancer prevention: reducing the risk of cancer with healthy food choices and physical activity". *Ca*. 56 (5): 254–81; quiz 313–4. doi:10.3322/canjclin.56.5.254. PMID 17005596.
40. Bhaskaran K, Douglas I, Forbes H, dos-Santos-Silva I, Leon DA, Smeeth L (August 2014). "Body-mass index and risk of 22 specific cancers: a population-based cohort study of 5·24 million UK adults". *Lancet*. 384 (9945): 755–65. doi:10.1016/S0140-6736(14)60892-8. PMID 25129328.
41. ^c Park S, Bae J, Nam BH, Yoo KY (2008). "Aetiology of cancer in Asia"(PDF). *Asian Pacific Journal of Cancer Prevention*. 9 (3): 371–80. PMID 18990005.
42. Brenner H, Rothenbacher D, Arndt V (2009). "Epidemiology of stomach cancer". *Methods in Molecular Biology*. *Methods in Molecular Biology*. 472: 467–77. doi:10.1007/978-1-60327-492-0_23. ISBN 978-1-60327-491-3. PMID 19107449.
43. Buell P, Dunn JE (May 1965). "CANCER MORTALITY AMONG JAPANESE ISSEI AND NISEI OF CALIFORNIA". *Cancer*. 18 (5): 656–64. doi:10.1002/1097-







0142(196505)18:5<656::AID-CNCR2820180515>3.0.CO;2-3. PMID 14278899.





44. Pagano JS, Blaser M, Buendia MA, Damania B, Khalili K, Raab-Traub N, Roizman B (December 2004). "Infectious agents and cancer: criteria for a causal relation". *Seminars in Cancer Biology*. 14 (6): 453–71. doi:10.1016/j.semcancer.2004.06.009. PMID 15489139.
45. Ljubojevic S, Skerlev M (2014). "HPV-associated diseases". *Clinics in Dermatology*. 32 (2): 227–34. doi:10.1016/j.clindermatol.2013.08.007. PMID 24559558.
46. Samaras V, Rafailidis PI, Mourtzoukou EG, Peppas G, Falagas ME (June 2010). "Chronic bacterial and parasitic infections and cancer: a review"(PDF). *Journal of Infection in Developing Countries*. 4 (5): 267–81. doi:10.3855/jidc.819. PMID 20539059.
47. Little JB (2000). "Chapter 14: Ionizing Radiation". In Kufe DW, Pollock RE, Weichselbaum RR, Bast RC, Gansler TS, Holland JF, Frei E. *Cancer medicine* (6th ed.). Hamilton, Ont: B.C. Decker. ISBN 1-55009-113-1.
48. Brenner DJ, Hall EJ (November 2007). "Computed tomography--an increasing source of radiation exposure". *The New England Journal of Medicine*. 357 (22):2277-84. doi:10.1056/NEJMra072149. PMID 18046031.
49. Cleaver JE, Mitchell DL (2000). "15. Ultraviolet Radiation Carcinogenesis". In Bast RC, Kufe DW, Pollock RE, et al. *Holland-Frei Cancer Medicine* (5th ed.). Hamilton, Ontario: B.C. Decker. ISBN 1-55009-113-1.
50. "IARC classifies radiofrequency electromagnetic fields as possibly carcinogenic to humans" (PDF). World Health Organization.
51. "Cell Phones and Cancer Risk - National Cancer Institute". Cancer.gov. 2013-05-08. Retrieved 2013-12-15.




52. Roukos DH (April 2009). "Genome-wide association studies: how predictable is a person's cancer risk?". *Expert Review of Anticancertherapy*. 9 (4):389–92. doi:10.1586/era.09.12. PMID 19374592.
53. Cunningham D, Atkin W, Lenz HJ, Lynch HT, Minsky B, Nordlinger B, Starling N (March 2010). "Colorectal cancer". *Lancet*. 375 (9719): 1030–47. doi:10.1016/S0140-6736(10)60353-4. PMID 20304247.
54. Maltoni CF, Holland JF (2000). "Chapter 16: Physical Carcinogens". In Bast RC, Kufe DW, Pollock RE, et al. *Holland-Frei Cancer Medicine* (5th ed.). Hamilton, Ontario: B.C. Decker. ISBN 1-55009-113-1. Retrieved 31 January 2011.
55. Gaeta, John F (2000). "Chapter 17: Trauma and Inflammation". In Bast RC, Kufe DW, Pollock RE, et al. *Holland-Frei Cancer Medicine* (5th ed.). Hamilton, Ontario: B.C. Decker. ISBN 1-55009-113-1. Retrieved 27 January 2011.
56. Colotta F, Allavena P, Sica A, Garlanda C, Mantovani A (July 2009). "Cancer-related inflammation, the seventh hallmark of cancer: links to genetic instability". *Carcinogenesis* (review). 30 (7): 1073–81. doi:10.1093/carcin/bgp127. PMID 19468060.
57. Ungefroren H, Sebens S, Seidl D, Lehnert H, Hass R (September 2011). "Interaction of tumor cells with the microenvironment". *Cell Communication and Signaling*. 9 (18): 18. doi:10.1186/1478-811X-9-18. PMID 21914164.
58. Mantovani A (June 2010). "Molecular pathways linking inflammation and cancer". *Current Molecular Medicine* (review). 10 (4): 369–73. doi:10.2174/156652410791316968. PMID 20455855.
59. Borrello MG, Degl'Innocenti D, Pierotti MA (August 2008). "Inflammation and cancer: the oncogene-driven


- connection". *Cancer Letters* (review). 267 (2): 262–70. doi:10.1016/j.canlet.2008.03.060. PMID 18502035.
60. Henderson BE, Bernstein L, Ross RK (2000). "Chapter 13: Hormones and the Etiology of Cancer". In Bast RC, Kufe DW, Pollock RE, et al. *Holland-Frei Cancer Medicine* (5th ed.). Hamilton, Ontario: B.C. Decker. ISBN 1-55009-113-1. Retrieved 27 January 2011.
 61. Rowlands MA, Gunnell D, Harris R, Vatten LJ, Holly JM, Martin RM (May 2009). "Circulating insulin-like growth factor peptides and prostate cancer risk: a systematic review and meta-analysis". *International Journal of Cancer*. 124 (10): 2416–29. doi:10.1002/ijc.24202. PMC 2743036  . PMID 19142965.
 62. Han Y, Chen W, Li P, Ye J (September 2015). "Association Between Coeliac Disease and Risk of Any Malignancy and Gastrointestinal Malignancy: A Meta-Analysis". *Medicine*. 94 (38): e1612. doi:10.1097/MD.0000000000001612. PMC 4635766  . PMID 26402826.
 63. Axelrad JE, Lichtiger S, Yajnik V (May 2016). "Inflammatory bowel disease and cancer: The role of inflammation, immunosuppression, and cancer treatment". *World Journal of Gastroenterology*. 22 (20): 4794–801. doi:10.3748/wjg.v22.i20.4794. PMC 4873872  . PMID 27239106.
 64. Croce CM (January 2008). "Oncogenes and cancer". *The New England Journal of Medicine*. 358 (5): 502–11. doi:10.1056/NEJMr072367. PMID 18234754.
 65. Knudson AG (November 2001). "Two genetic hits (more or less) to cancer". *Nature Reviews. Cancer*. 1 (2): 157–62. doi:10.1038/35101031. PMID 11905807.
 66. Nelson DA, Tan TT, Rabson AB, Anderson D, Degenhardt K, White E (September 2004). "Hypoxia and defective apoptosis drive genomic instability and

- tumorigenesis". *Genes & Development*. 18 (17): 2095–107. doi:10.1101/gad.1204904. PMC 515288  . PMID 15314031.
67. Merlo LM, Pepper JW, Reid BJ, Maley CC (December 2006). "Cancer as an evolutionary and ecological process". *Nature Reviews. Cancer*. 6 (12): 924–35. doi:10.1038/nrc2013. PMID 17109012.
 68. Baylin SB, Ohm JE (February 2006). "Epigenetic gene silencing in cancer - a mechanism for early oncogenic pathway addiction?". *Nature Reviews. Cancer*. 6 (2): 107–16. doi:10.1038/nrc1799. PMID 16491070.
 69. Kanwal R, Gupta S (April 2012). "Epigenetic modifications in cancer". *Clinical Genetics*. 81 (4): 303–11. doi:10.1111/j.1399-0004.2011.01809.x. PMC 3590802  . PMID 22082348.
 70. Baldassarre G, Battista S, Belletti B, Thakur S, Pentimalli F, Trapasso F, Fedele M, Pierantoni G, Croce CM, Fusco A (April 2003). "Negative regulation of BRCA1 gene expression by HMGA1 proteins accounts for the reduced BRCA1 protein levels in sporadic breast carcinoma". *Molecular and Cellular Biology*. 23 (7): 2225–38. doi:10.1128/MCB.23.7.2225-2238.2003. PMC 150734  . PMID 12640109./
 71. Schnekenburger M, Diederich M (March 2012). "Epigenetics Offer New Horizons for Colorectal Cancer Prevention". *Current Colorectal Cancer Reports*. 8 (1): 66–81. doi:10.1007/s11888-011-0116-z. PMC 3277709  . PMID 22389639.
 72. Jacinto FV, Esteller M (July 2007). "Mutator pathways unleashed by epigenetic silencing in human cancer". *Mutagenesis*. 22 (4): 247–53. doi:10.1093/mutage/gem009. PMID 17412712.
 73. Lahtz C, Pfeifer GP (February 2011). "Epigenetic changes of DNA repair genes in cancer". *Journal of Molecular*


8. doi:10.1093/jmcb/mjq053. PMC 3030973  . PMID 21278452.
74. Bernstein C, Nfonsam V, Prasad AR, Bernstein H (March 2013). "Epigenetic field defects in progression to cancer". *World Journal of Gastrointestinal Oncology*. 5 (3): 43–9. doi:10.4251/wjgo.v5.i3.43. PMC 3648662  . PMID 23671730.
75. Bernstein C, Prasad AR, Nfonsam V, Bernstein H (2013). "DNA Damage, DNA Repair and Cancer". In Chen C. *New Research Directions in DNA Repair*. InTech. doi:10.5772/53919. ISBN 978-953-51-1114-6.
76. Narayanan L, Fritzell JA, Baker SM, Liskay RM, Glazer PM (April 1997). "Elevated levels of mutation in multiple tissues of mice deficient in the DNA mismatch repair gene Pms2". *Proceedings of the National Academy of Sciences of the United States of America*. 94 (7): 3122–7. doi:10.1073/pnas.94.7.3122. PMC 20332  . PMID 9096356.
77. Hegan DC, Narayanan L, Jirik FR, Edelmann W, Liskay RM, Glazer PM (December 2006). "Differing patterns of genetic instability in mice deficient in the mismatch repair genes Pms2, Mlh1, Msh2, Msh3 and Msh6". *Carcinogenesis*. 27 (12): 2402–8. doi:10.1093/carcin/bgl079. PMC 2612936  . PMID 16728433.
78. Tutt AN, van Oostrom CT, Ross GM, van Steeg H, Ashworth A (March 2002). "Disruption of Brca2 increases the spontaneous mutation rate in vivo: synergism with ionizing radiation". *EMBO Reports*. 3 (3): 255–60. doi:10.1093/embo-reports/kvf037. PMC 1084010  . PMID 11850397.
79. German J (March 1969). "Bloom's syndrome. I. Genetical and clinical observations in the first twenty-seven patients". *American Journal of Human Genetics*. 21 (2): 196–227. PMC 1706430  . PMID 5770175.

80. O'Hagan HM, Mohammad HP, Baylin SB (August 2008). Lee JT, ed. "Double strand breaks can initiate gene silencing and SIRT1-dependent onset of DNA methylation in an exogenous promoter CpG island". *PLoS Genetics*. 4 (8): e1000155. doi:10.1371/journal.pgen.1000155. PMC 2491723 . PMID 18704159.
81. Cuozzo C, Porcellini A, Angrisano T, Morano A, Lee B, Di Pardo A, Messina S, Iuliano R, Fusco A, Santillo MR, Muller MT, Chiariotti L, Gottesman ME, Avvedimento EV (July 2007). "DNA damage, homology-directed repair, and DNA methylation". *PLoS Genetics*. 3 (7): e110. doi:10.1371/journal.pgen.0030110. PMC 1913100 . PMID 17616978.
82. Malkin D (April 2011). "Li-fraumeni syndrome". *Genes & Cancer*. 2 (4):475–doi:10.1177/1947601911413466. PMC 3135649 . PMID 21779515.
83. Fearon ER (November 1997). "Human cancer syndromes: clues to the origin and nature of cancer". *Science*. 278 (5340): 1043–50. doi:10.1126/science.278.5340.1043. PMID 9353177.
84. Vogelstein B, Papadopoulos N, Velculescu VE, Zhou S, Diaz LA, Kinzler KW (March 2013). "Cancer genome landscapes". *Science*. 339 (6127): 1546–58. doi:10.1126/science.1235122. PMC 3749880 . PMID 23539594.
85. "Metastatic Cancer: Questions and Answers". National Cancer Institute. Retrieved 2008-08-28.
86. "What is Metastasized Cancer?". National Comprehensive Cancer Network. Archived from the original on 7 July 2013. Retrieved 18 July 2013.
87. Varricchio Claudette G. (2004). *A cancer source book for nurses*. Boston: Jones and Bartlett Publishers. p. 229. ISBN 0-7637-3276-1.

88. "Cancer prevention: 7 steps to reduce your risk". 2008 Mayo Clinic. 27 September 2008. Retrieved 30 January 2010.
89. Danaei G, Vander Hoorn S, Lopez AD, Murray CJ, Ezzati M (November 2005). "Causes of cancer in the world: comparative risk assessment of nine behavioural and environmental risk factors". *Lancet*. 366 (9499): 1784–93. doi:10.1016/S0140-6736(05)67725-2. PMID 16298215.
90. Wu S, Powers S, Zhu W, Hannun YA (January 2016). "Substantial contribution of extrinsic risk factors to cancer development". *Nature*. 529 (7584): 43–7. doi:10.1038/nature16166. PMC 4836858 . PMID 26675728.
91. "Cancer". World Health Organization. Retrieved 9 January 2011.
92. Wicki A, Hagmann J (September 2011). "Diet and cancer". *Swiss Medical Weekly*. 141: w13250. doi:10.4414/smw.2011.13250. PMID 21904992.
93. Cappellani A, Di Vita M, Zanghi A, Cavallaro A, Piccolo G, Veroux M, Berretta M, Malaguarnera M, Canzonieri V, Lo Menzo E (January 2012). "Diet, obesity and breast cancer: an update". *Frontiers in Bioscience*. 4: 90–108. PMID 22202045.
94. Key TJ (January 2011). "Fruit and vegetables and cancer risk". *British Journal of Cancer*. 104 (1): 6–11. doi:10.1038/sj.bjc.6606032. PMC 3039795 . PMID 21119663. Wang X, Ouyang Y, Liu J, Zhu M, Zhao G, Bao W, Hu FB (July 2014). "Fruit and vegetable consumption and mortality from all causes, cardiovascular disease, and cancer: systematic review and dose-response meta-analysis of prospective cohort studies". *Bmj*. 349: g4490. doi:10.1136/bmj.g4490. PMC 4115152 . PMID 25073782.
95. Larsson SC, Wolk A (May 2007). "Coffee consumption and risk of liver cancer: a meta-

- analysis". *Gastroenterology*. 132 (5): 1740–5. doi:10.1053/j.gastro.2007.03.044. PMID 17484871.
96. Zheng W, Lee SA (2009). "Well-done meat intake, heterocyclic amine exposure, and cancer risk". *Nutrition and Cancer*. 61 (4): 437–46. doi:10.1080/01635580802710741. PMC 2769029  . PMID 19838915.
97. Ferguson LR (February 2010). "Meat and cancer". *Meat Science*. 84 (2): 308–13. doi:10.1016/j.meatsci.2009.06.032. PMID 20374790.
98. Staff (October 26, 2015). "World Health Organization - IARC Monographs evaluate consumption of red meat and processed meat" (PDF). International Agency for Research on Cancer. Retrieved October 26, 2015.
99. Hauser, Christine (October 26, 2015). "W.H.O. Report Links Some Cancers With Processed or Red Meat". *New York Times*. Retrieved October 26, 2015.
100. Holland Chp.33
101. Rostom A, Dubé C, Lewin G, Tsertsvadze A, Barrowman N, Code C, Sampson M, Moher D (March 2007). "Nonsteroidal anti-inflammatory drugs and cyclooxygenase-2 inhibitors for primary prevention of colorectal cancer: a systematic review prepared for the U.S. Preventive Services Task Force". *Annals of Internal Medicine*. 146 (5): 376–89. doi:10.7326/0003-4819-146-5-200703060-00010. PMID 17339623.
102. Rothwell PM, Fowkes FG, Belch JF, Ogawa H, Warlow CP, Meade TW (January 2011). "Effect of daily aspirin on long-term risk of death due to cancer: analysis of individual patient data from randomised trials". *Lancet*. 377 (9759): 31–41. doi:10.1016/S0140-6736(10)62110-1. PMID 21144578.
103. Cooper K, Squires H, Carroll C, Papaioannou D, Booth A, Logan RF, Maguire C, Hind D, Tappenden P (June 2010). "Chemoprevention of colorectal cancer: systematic review and economic evaluation". *Health Technology*

- Assessment. 14 (32): 1–206. doi:10.3310/hta14320. PMID 20594533.
104. Thomsen A, Kolesar JM (December 2008). "Chemoprevention of breast cancer". *American Journal of Health-System Pharmacy*. 65 (23): 2221–8. doi:10.2146/ajhp070663. PMID 19020189.
 105. Wilt TJ, MacDonald R, Hagerty K, Schellhammer P, Kramer BS (April 2008). Wilt TJ, ed. "Five-alpha-reductase Inhibitors for prostate cancer prevention". *The Cochrane Database of Systematic Reviews* (2): CD007091. doi:10.1002/14651858.CD007091. PMID 18425978.
 106. "Vitamins and minerals: not for cancer or cardiovascular prevention". *Prescrire International*. 19 (108): 182. August 2010. PMID 20939459.
 107. Giovannucci E, Liu Y, Rimm EB, Hollis BW, Fuchs CS, Stampfer MJ, Willett WC (April 2006). "Prospective study of predictors of vitamin D status and cancer incidence and mortality in men". *Journal of the National Cancer Institute*. 98 (7): 451–9. doi:10.1093/jnci/djj101. PMID 16595781.
 108. "Vitamin D Has Role in Colon Cancer Prevention". Archived from the original on 4 December 2006. Retrieved 27 July 2007.
 109. Holick MF (January 2013). "Vitamin D, sunlight and cancer connection". *Anti-Cancer Agents in Medicinal Chemistry*. 13 (1): 70–82. doi:10.2174/187152013804487308. PMID 23094923.
 110. Schwartz GG, Blot WJ (April 2006). "Vitamin D status and cancer incidence and mortality: something new under the sun". *Journal of the National Cancer Institute*. 98 (7): 428–30. doi:10.1093/jnci/djj127. PMID 16595770.
 111. Fritz H, Kennedy D, Fergusson D, Fernandes R, Doucette S, Cooley K, Seely A, Sagar S, Wong R, Seely D

- (2011). Minna JD, ed. "Vitamin A and retinoid derivatives for lung cancer: a systematic review and meta analysis". PloS One. 6 (6): e21107. Bibcode:2011PLoSO...6E1107F. doi:10.1371/journal.pone.0021107. PMC 3124481 . PMID 21738614.
112. Cole BF, Baron JA, Sandler RS, Haile RW, Ahnen DJ, Bresalier RS, McKeown-Eyssen G, Summers RW, Rothstein RI, Burke CA, Snover DC, Church TR, Allen JI, Robertson DJ, Beck GJ, Bond JH, Byers T, Mandel JS, Mott LA, Pearson LH, Barry EL, Rees JR, Marcon N, Saibil F, Ueland PM, Greenberg ER (June 2007). "Folic acid for the prevention of colorectal adenomas: a randomized clinical trial". *Jama*. 297 (21): 2351–9. doi:10.1001/jama.297.21.2351. PMID 17551129.
 113. Vinceti M, Dennert G, Crespi CM, Zwahlen M, Brinkman M, Zeegers MP, Horneber M, D'Amico R, Del Giovane C (March 2014). "Selenium for preventing cancer". *The Cochrane Database of Systematic Reviews*. 3 (3): CD005195. doi:10.1002/14651858.CD005195.pub3. PMID 24683040.
 114. "Cancer Vaccine Fact Sheet". NCI. 8 June 2006. Retrieved 15 November 2008.
 115. Lertkhachonsuk AA, Yip CH, Khuhaprema T, Chen DS, Plummer M, Jee SH, Toi M, Wilailak S (November 2013). "Cancer prevention in Asia: resource-stratified guidelines from the Asian Oncology Summit 2013". *The Lancet. Oncology*. 14 (12): e497–507. doi:10.1016/S1470-2045(13)70350-4. PMID 24176569.
 116. Wilson JMG, Jungner G. (1968) Principles and practice of screening for disease. Geneva:World Health Organization. Public Health Papers, #34.
 117. "Screening for Cervical Cancer". U.S. Preventive Services Task Force. 2003.

118. Dong-Jiann HUANG¹, Chun-Der LIN¹, Hsien-Jung CHEN², and Yaw-Huei LIN^{1,*} 2004 Antioxidant and antiproliferative activities of sweet potato (*Ipomoea batatas* [L.] Lam 'Tainong 57') constituents Bot. Bull. Acad. Sin. 45: 179-186
119. Prasanth nv*, dilip c, sanal dev kt, lis augustine, saraswathi r. 2010 evaluation of invitro cytotoxic and antioxidant activities of ipomoea batatas International Journal of Pharmacy and Pharmaceutical Sciences ISSN- 0975-1491 Vol 2, Issue 3
120. Seow-Mun Hue, Amru Nasrulhaq Boyce, Chandran Somasundram 2011 Comparative Study on the Antioxidant Activity of Leaf Extract and Carotenoids Extract from *Ipomoea batatas* var. Oren (Sweetpotato) Leaves International Journal of Biological, Biomolecular, Agricultural, Food and Biotechnological Engineering Vol:5, No:10
121. Vandana Panda¹ , Madhav Sonkamble¹ , and Swati Patil² 2011 Wound healing activity of *Ipomoea batatas* tubers (sweet potato) Functional Foods in Health and Disease 10:403-415
122. Vandana Panda* and Madhav Sonkamble 2012 Anti-ulcer activity of *Ipomoea batatas* tubers (sweet potato) Functional Foods in Health and Disease 2(3):48-61
123. V.B. Adsull*, E.Khathiwora² , Rasika Torane² , N.R. Deshpande 2012 Antimicrobial activities of *Ipomoea carnea* leaves J. Nat. Prod. Plant Resour., 2 (5):597-600
124. Marilena Meira,^{*},^{1*} Eliezer Pereira da Silva,² Jorge M. David,^{Juceni P. david}³ 2012 Review of the genus *Ipomoea*: traditional uses, chemistry and biological activities Revista Brasileira de Farmacognosia Brazilian Journal of Pharmacognosy 22(3): 682-713, May/Jun. 2012

125. G. Anand*, G. Sumithira, R. Chinna Raja, A. Muthukumar, G. Vidhya 2013 In vitro and in vivo anticancer activity of hydroalcoholic extract of Ipomoea arnea leaf against Ehrlich Ascites Carcinoma cell lines Anand G et al; Int J Adv Pharm Gen Res, 1(1), 39-54
126. Yuzhi Jiao¹, Yanjie Jiang², Weiwei Zhai¹ and Zhendong Yang^{1*} 2012 Studies on antioxidant capacity of anthocyanin extract from purple sweet potato (*Ipomoea batatas* L.) African Journal of Biotechnology Vol. 11(27), pp. 7046-7054
127. Raju Asirvatham^{1*}, Arockiasamy Josphin Maria Christina², Anita Murali³ 2013 In Vitro Antioxidant and Anticancer Activity Studies on *Drosera Indica* L. (*Droseraceae*) Advanced Pharmaceutical Bulletin, 3(1), 115-120
128. Eleazu C. O.^{1*} and Ironua C.² 2013 february Physicochemical composition and antioxidant properties of a sweetpotato variety (*Ipomoea batatas* L) commercially sold in South Eastern Nigeria African Journal of Biotechnology Vol. 12(7), pp. 720-727
129. Zainal Baharum , Abdah Md Akim , Yun Hin Taufiq-Yap , Roslida Abdul Hamid and Rosmin Kasran 2014 In Vitro Antioxidant and Antiproliferative Activities of Methanolic Plant Part Extracts of *Theobroma cacao* Molecules 2014, 19, 18317-18331; doi:10.3390/molecules191118317
130. Marcelia Sugata, Chien-Yih Lin, and Yang-Chia Shih 2015 Anti-Inflammatory and Anticancer Activities of Taiwanese Purple-Fleshed Sweet Potatoes (*Ipomoea batatas* L. Lam) Extracts
131. Parle Milind and Monika 2015 sweet potato as super food. Parle Milind et.al/ Ind. J. Res. Ayurvedha pharm. 6(4)
132. BioMed Research International Volume 2015, Article ID 768093, 10 pages

133. M. Sucharitha¹, M. Kotes^{*1}, K. Devika², Y. Naresh³,
M. Kiran⁴ 2016 Evaluation of Diuretic Activity of aqueous
extract of Ipomoea batatas (L) Sch. J. App. Med. Sci., 2016;
4(6A):1902-1905